

New prime K-tuple theorem (6)

$$P, P+4^n (n=1, \dots, k)$$

Jiang, Chunxuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: Using Jiang function we prove that for every positive integer k there exist infinitely many primes P such that each of $P+4^n$ is prime.

[Chun-Xuan Jiang. New prime K-tuple theorem (6) $P, P+4^n (n=1, \dots, k)$. Academ Arena 2016;8(2s): 9-9]. (ISSN 1553-992X). <http://www.sciencepub.net/academia>. 6. doi:[10.7537/marsaj0802s1606](https://doi.org/10.7537/marsaj0802s1606).

Keywords: new; prime; k -tuple; theorem; Jiang Chunxuan; mathematics; science; number; function

Theorem

$$P, P+4^n (n=1, \dots, k) \quad (1)$$

For every positive integer k there exist infinitely many primes P such that each of $P+4^n$ is prime.

Proof. We have Jiang function [1]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)], \quad (2)$$

$$\omega = \prod_P P,$$

$\chi(P)$ is the number of solutions of congruence

$$\prod_{n=1}^k [q + 4^n] \equiv 0 \pmod{P}, \quad (3)$$

where $q = 1, \dots, P-1$.

From (3) we have

$$\text{If } P < 2k \text{ then } \chi(P) = \frac{P-1}{2}, \text{ if } 2k < P \text{ then } \chi(P) = k.$$

From (3) and (2) we have

$$J_3(\omega) = \prod_{P=3}^{P<2k} \frac{P-1}{2} \prod_{2k < P} (P-1-k) \neq 0 \quad (4)$$

We prove that for every positive integer k there exist infinitely many primes p such that each of $P+4^n$ is prime.
We have the best asymptotic formula [1, 2]

$$\pi_{k+1}(N, 2) = \left| \left\{ P \leq N : P+4^n = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N} \quad (5)$$

$$\text{where } \phi(\omega) = \prod_P (P-1)$$

References

Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>)