

## The Hardy-Littlewood prime $k$ -tuple conjecture is false

Jiang, Chun-Xuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

[jiangchunxuan@sohu.com](mailto:jiangchunxuan@sohu.com), [cxjiang@mail.bcf.net.cn](mailto:cxjiang@mail.bcf.net.cn), [jcxuan@sina.com](mailto:jcxuan@sina.com), [Jiangchunxuan@vip.sohu.com](mailto:Jiangchunxuan@vip.sohu.com),  
[jcxxxx@163.com](mailto:jcxxxx@163.com)

**Abstract:** Using Jiang function we prove Jiang prime  $k$ -tuple theorem. We prove that the Hardy-Littlewood prime  $k$ -tuple conjecture is false. Jiang prime  $k$ -tuple theorem can replace the Hardy-Littlewood prime  $k$ -tuple conjecture.

[Jiang, Chun-Xuan (蒋春暄). **The Hardy-Littlewood prime  $k$ -tuple conjecture is false.** *Academ Arena* 2016;8(6):44-48]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 9. doi:[10.7537/marsaa080616.09](https://doi.org/10.7537/marsaa080616.09).

**Keywords:** Jiang function; prime  $k$ -tuple theorem; Hardy-Littlewood prime;  $k$ -tuple conjecture

### (A) Jiang prime $k$ -tuple theorem [1, 2].

We define the prime  $k$ -tuple equation

$$p, p + n_i, \dots, (1)$$

where  $2|n_i, i = 1, \dots, k-1$

we have Jiang function [1, 2]

$$J_2(\omega) = \prod_P (P-1 - \chi(P)), \quad (2)$$

where  $\omega = \prod_P P$ ,  $\chi(P)$  is the number of solutions of congruence

$$\prod_{i=1}^{k-1} (q + n_i) \equiv 0 \pmod{P}, \quad q = 1, \dots, p-1. \quad (3)$$

If  $\chi(P) < P-1$  then  $J_2(\omega) \neq 0$ . There exist infinitely many primes  $P$  such that each of  $P + n_i$  is prime. If  $\chi(P) = P-1$  then  $J_2(\omega) = 0$ . There exist finitely many primes  $P$  such that each of  $P + n_i$  is prime.  $J_2(\omega)$  is a subset of Euler function  $\phi(\omega)$  [2].

If  $J_2(\omega) \neq 0$ , then we have the best asymptotic formula of the number of prime  $P$  [1, 2]

$$\pi_k(N, 2) = |\{P \leq N : P + n_i = \text{prime}\}| \sim \frac{J_2(\omega) \omega^{k-1}}{\phi^k(\omega)} \frac{N}{\log^k N} = C(k) \frac{N}{\log^k N} \quad (4)$$

$$\phi(\omega) = \prod_P (P-1),$$

$$C(k) = \prod_P \left(1 - \frac{1 + \chi(P)}{P}\right) \left(1 - \frac{1}{P}\right)^{-k} \quad (5)$$

**Example 1.** Let  $k = 2, P, P+2$ , twin primes theorem.

From (3) we have

$$\chi(2) = 0, \quad \chi(P) = 1 \quad \text{if } P > 2, \quad (6)$$

Substituting (6) into (2) we have

$$J_2(\omega) = \prod_{P \geq 3} (P - 2) \neq 0 \quad (7)$$

There exist infinitely many primes  $P$  such that  $P + 2$  is prime. Substituting (7) into (4) we have the best asymptotic formula

$$\pi_k(N, 2) = |\{P \leq N : P + 2 = \text{prime}\}| \sim 2 \prod_{P \geq 3} \left(1 - \frac{1}{(P-1)^2}\right) \frac{N}{\log^2 N}. \quad (8)$$

**Example 2.** Let  $k = 3, P, P + 2, P + 4$ .

From (3) we have

$$\chi(2) = 0, \quad \chi(3) = 2 \quad (9)$$

From (2) we have

$$J_2(\omega) = 0 \quad (10)$$

It has only a solution  $P = 3, P + 2 = 5, P + 4 = 7$ . One of  $P, P + 2, P + 4$  is always divisible by 3.

**Example 3.** Let  $k = 4, P, P + n$ , where  $n = 2, 6, 8$ .

From (3) we have

$$\chi(2) = 0, \quad \chi(3) = 1, \quad \chi(P) = 3 \quad \text{if } P > 3 \quad (11)$$

Substituting (11) into (2) we have

$$J_2(\omega) = \prod_{P \geq 5} (P - 4) \neq 0 \quad (12)$$

There exist infinitely many primes  $P$  such that each of  $P + n$  is prime.

Substituting (12) into (4) we have the best asymptotic formula

$$\pi_4(N, 2) = |\{P \leq N : P + n = \text{prime}\}| \sim \frac{27}{3} \prod_{P \geq 5} \frac{P^3(P-4)}{(P-1)^4} \frac{N}{\log^4 N} \quad (13)$$

**Example 4.** Let  $k = 5, P, P + n$ , where  $n = 2, 6, 8, 12$ .

From (3) we have

$$\chi(2) = 0, \quad \chi(3) = 1, \quad \chi(5) = 3, \quad \chi(P) = 4 \quad \text{if } P > 5 \quad (14)$$

Substituting (14) into (2) we have

$$J_2(\omega) = \prod_{P \geq 7} (P - 5) \neq 0 \quad (15)$$

There exist infinitely many primes  $P$  such that each of  $P + n$  is prime. Substituting (15) into (4) we have the best asymptotic formula

$$\pi_5(N, 2) = |\{P \leq N : P + n = \text{prime}\}| \sim \frac{15^4}{2^{11}} \prod_{P \geq 7} \frac{(P-5)P^4}{(P-1)^5} \frac{N}{\log^5 N} \quad (16)$$

**Example 5.** Let  $k = 6, P, P + n$ , where  $n = 2, 6, 8, 12, 14$ .

From (3) and (2) we have

$$\chi(2) = 0, \quad \chi(3) = 1, \quad \chi(5) = 4, \quad J_2(5) = 0 \quad (17)$$

It has only a solution  $P = 5, P + 2 = 7, P + 6 = 11, P + 8 = 13, P + 12 = 17, P + 14 = 19$ . One of  $P + n$  is always divisible by 5.

**(B) The Hardy-Littlewood prime  $k$ -tuple conjecture[3-14].**

This conjecture is generally believed to be true, but has not been proved(Odlyzko et al.1999).

We define the prime  $k$ -tuple equation

$$P, P+n_i \quad (18)$$

where  $2|n_i, i=1, \dots, k-1$

In 1923 Hardy and Littlewood conjectured the asymptotic formula

$$\pi_k(N, 2) = |\{P \leq N : P+n_i = \text{prime}\}| \sim H(k) \frac{N}{\log^k N}, \quad (19)$$

where

$$H(k) = \prod_P \left(1 - \frac{\nu(P)}{P}\right) \left(1 - \frac{1}{P}\right)^{-k} \quad (20)$$

$\nu(P)$  is the number of solutions of congruence

$$\prod_{i=1}^{k-1} (q+n_i) \equiv 0 \pmod{P}, \quad q=1, \dots, P \quad (21)$$

From (21) we have  $\nu(P) < P$  and  $H(k) \neq 0$ . For any prime  $k$ -tuple equation there exist infinitely many primes  $P$  such that each of  $P+n_i$  is prime, which is false.

**Conjecture 1.** Let  $k=2, P, P+2$ , twin primes theorem

From (21) we have

$$\nu(P)=1 \quad (22)$$

Substituting (22) into (20) we have

$$H(2) = \prod_P \frac{P}{P-1} \quad (23)$$

Substituting (23) into (19) we have the asymptotic formula

$$\pi_2(N, 2) = |\{P \leq N : P+2 = \text{prime}\}| \sim \prod_P \frac{P}{P-1} \frac{N}{\log^2 N} \quad (24)$$

which is false see example 1.

**Conjecture 2.** Let  $k=3, P, P+2, P+4$ .

From (21) we have

$$\nu(2)=1, \nu(P)=2 \quad \text{if } P>2 \quad (25)$$

Substituting (25) into (20) we have

$$H(3) = 4 \prod_{P \geq 3} \frac{P^2(P-2)}{(P-1)^3} \quad (26)$$

Substituting (26) into (19) we have asymptotic formula

$$\pi_3(N, 2) = |\{P \leq N : P+2 = \text{prime}, P+4 = \text{prime}\}| \sim 4 \prod_{P \geq 3} \frac{P^2(P-2)}{(P-1)^3} \frac{N}{\log^3 N} \quad (27)$$

which is false see example 2.

**Conjecture 3.** Let  $k = 4$ ,  $P, P+n$ , where  $n = 2, 6, 8$ .

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(P) = 3 \text{ if } P > 3 \quad (28)$$

Substituting (28) into (20) we have

$$H(4) = \frac{27}{2} \prod_{P>3} \frac{P^3(P-3)}{(P-1)^4} \quad (29)$$

Substituting (29) into (19) we have asymptotic formula

$$\pi_4(N, 2) = |\{P \leq N : P+n = \text{prime}\}| \sim \frac{27}{2} \prod_{P>3} \frac{P^3(P-3)}{(P-1)^4} \frac{N}{\log^4 N} \quad (30)$$

Which is false see example 3.

**Conjecture 4.** Let  $k = 5$ ,  $P, P+n$ , where  $n = 2, 6, 8, 12$

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(5) = 3, \nu(P) = 4 \text{ if } P > 5 \quad (31)$$

Substituting (31) into (20) we have

$$H(5) = \frac{15^4}{4^5} \prod_{P>5} \frac{P^4(P-4)}{(P-1)^5} \quad (32)$$

Substituting (32) into (19) we have asymptotic formula

$$\pi_5(N, 2) = |\{P \leq N : P+n = \text{prime}\}| \sim \frac{15^4}{4^5} \prod_{P>5} \frac{P^4(P-4)}{(P-1)^5} \frac{N}{\log^5 N} \quad (33)$$

Which is false see example 4.

**Conjecture 5.** Let  $k = 6$ ,  $P, P+n$ , where  $n = 2, 6, 8, 12, 14$ .

From (21) we have

$$\nu(2) = 1, \nu(3) = 2, \nu(5) = 4, \nu(P) = 5 \text{ if } P > 5 \quad (34)$$

Substituting (34) into (20) we have

$$H(6) = \frac{15^5}{2^{13}} \prod_{P>5} \frac{(P-5)P^5}{(P-1)^6} \quad (35)$$

Substituting (35) into (19) we have asymptotic formula

$$\pi_6(N, 2) = |\{P \leq N : P+n = \text{prime}\}| \sim \frac{15^5}{2^{13}} \prod_{P>5} \frac{(P-5)P^5}{(P-1)^6} \frac{N}{\log^6 N} \quad (36)$$

which is false see example 5.

## Conclusion.

The Hardy-Littlewood prime  $k$ -tuple conjecture is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime  $k$ -tuple conjecture. Jiang prime  $k$ -tuple theorem can replace Hardy-Littlewood prime  $k$ -tuple Conjecture. There cannot be really modern prime theory without Jiang function.

## References

1. Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press, 2002, MR2004c: 11001 (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>).
2. Chun-Xuan Jiang, Jiang's function  $J_{n+1}(\omega)$  in prime distribution. (<http://www.wbabin.net/math/xuan2.pdf>)

- 
- 3. (http://vixra.org/pdf/0812.0004v2.pdf).
  - 4. G. H. Hardy and J. E. Littlewood, Some problems of ‘Partition Numerorum’, III: On the expression of a number as a sum of primes, *Acta Math.*, 44(1923), 1-70.
  - 4. B. Green and T. Tao, The primes contain arbitrarily long arithmetic progressions, *Ann. Math.*, 167(2008), 481-547.
  - 5. D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yıldırım, Small gaps between products of two primes, *Proc. London Math. Soc.*, (3) 98 (2009) 741-774.
  - 6. D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yıldırım, Small gaps between primes or almost primes, *Trans. Amer. Math. Soc.*, 361(2009) 5285-5330.
  - 7. D. A. Goldston, J. Pintz and C. Y. Yıldırım, Primes in tuples I, *Ann. Math.*, 170(2009) 819-862.
  - 8. P. Ribenboim, The new book of prime number records, 3rd edition, Springer-Verlag, New York, NY, 1995. PP409-411.
  - 9. H. Halberstam and H.-E. Richert, Sieve methods, Academic Press, 1974.
  - 10. A. Schinzel and W. Sierpinski, Sur certaines hypothèses concernant les nombres premiers, *Acta Arith.*, 4(1958)185-208.
  - 11. P.T. Bateman and R.A. Horn, A heuristic asymptotic formula concerning the distribution of prime numbers, *Math.Comp.*,16(1962)363-367.
  - 12. W. Narkiewicz, The development of prime number theory, From Euclid to Hardy and Littlewood, Springer-Verlag, New York, NY, 2000, 333-53.
  - 13. B. Green and T. Tao, Linear equations in primes, To appear, *Ann. Math.*.
  - 14. T. Tao, Recent progress in additive prime number theory,  
<http://terrytao.files.wordpress.com/2009/08/prime-number-theory1.pdf>.
  - 15. Ma H. The Nature of Time and Space. *Nature and science* 2003;1(1):1-11. *Nature and science* 2007;5(1):81-96.

6/22/2016