

The role of endurance exercise program in hyperthyroidism in relation to ACE genotype

Heshmat, H.* and Saad El Taib**

* Dep. of physiol. Fac. Of Med. Zagazig Univ.

** Dep. of Biol. SC. and Motion Rehab. Fac. Of PE, Zawia Univ.

Taibsaad2010@yahoo.com; Husseinheshmat43@hotmail.com

Abstract: Background: There are three widely recognized types of exercise: endurance, resistance and sprint. Endurance exercise is characterized by prolonged continuous or intermittently periods of contractile activity against low resistance. ACE gene polymorphisms has been associated with some metabolic disorders. **Purpose :** This study investigated the role of endurance exercise program in hyperthyroidism in relation to ACE genotype. **Methods :** Using PCR method the ACE was genotyped in ten hyperthyroid patients and ten control before and after 12 weeks of endurance exercise program, thyroid hormones and TSH were investigated using Elisa technique. **Results :** ACEDD (80%) and ID (20%) genotype were associated with hyperthyroidism, in control, ACEID (50%), ACEII (25%), ACEDD (25%). TSH decreased in hyperthyroidism with increased F. T₄, T₃, T. T₄, T₃. compared to control. After endurance training program TSH increased, while F.T₄, T₃ and TT₄, T₃ decreased in hyperthyroidism together with reduction in pulse rate and blood pressure. **Conclusion:** ACE genotype may have a pathogenic role in thyroid gland, Endurance exercise training might have a positive effect alone in treating hyperthyroid subjects. **Recommendation:** to use ACE genotype to evaluate hyperthyroidism, and TSH and thyroid hormones as indicators of the efficiency of endurance training program.

[Heshmat, H. and Saad El Taib. **The role of endurance exercise program in hyperthyroidism in relation to ACE genotype.** *J Am Sci* 2013;9(2):456 -460]. (ISSN: 1545-1003). <http://www.jofamericanscience.org>. 63

1. Introduction

Thyroid hormones are essential for normal growth, development and metabolism. Hormone production by the thyroid gland is tightly regulated through the hypothalamic pituitary axis (Smith et al., 1990). They also added that thyroid disease is common, particularly in women, with a prevalence in the community of 3-5Y. Once diagnosed, thyroid disease is easily treated, with an excellent long-term outcome for most patients.

Guyton and Hall (2006) reported that the most important regulator of thyroid homeostasis is thyroid-stimulating hormone (TSH). This peptide hormone comprises a specific beta unit, which is required for binding to the TSH receptor, and an alpha subunit, which is common to the gonadotrophins, both subunits are required for bioactivity.

Hyperthyroidism is the symptom of an overactive thyroid gland follow logically from the actions of thyroid hormone, it is characterized by nervousness, weight loss, hyperphagia, heat intolerance, increased pulse pressure, a fine tremor of the fingers, sweating and a BMR from +10 to as high as + 100. The most common cause is graves disease, which accounts for 60 – 80% of the cases, This is an autoimmune disease, more common in women, in which antibodies to the TSH receptor stimulate the receptor. This produces marked T₄ and T₃ and enlargement of the thyroid gland (goiter), TSH is low and exophthalmos occurs in 50% of patients (Barret et al., 2010).

The renin – angiotensin – aldosterone system: Renin is secreted in response to a fall in renal afferent arteriolar pressure or to a reduction in supply of Na⁺ to the distal tubule. It convert angiotensinogen in plasma to angiotensin I (A_I), which in turn is converted to angiotensin II (A_{II}) by angiotensin converting enzyme (ACE). Both A_{II} and its metabolic product angiotensin III are pharmacologically active, and stimulate the release of aldosterone from the adrenal cortex, which promote Na⁺ reabsorption in exchange for urinary loss of H⁺ or K⁺ (El Shafei et al, 2011, Joshi et al 2011). Palvic et al., (2012) stated that ACE is an important regulator of blood pressure and cardiovascular homeostasis. Plasma levels of ACE depend on an insertion / deletion polymorphism in its gene.

El Taib (2012) reported that the thyroid gland hormones are so vital for many physiological process in the human body but the biological effects of short term changes on the levels of these hormones due to physical effort is not totally understood until now. Some studies mentioned that physical efforts lead to an instant decline in T₃ and T₄ concentrations, where others mentioned that physical effort has no effect on these hormones. A third group mentioned that physical efforts raised the level of these hormones as a reaction of exercises. The disagreement between those studies leads to the necessary of performing further researches trying to indentify this effect together with its action on ACE genotype.

Sports activity cause basic changes in the energy required for metabolism and to ensure the

excess production in muscle and cardiac contraction due to physical exertion and the hormones working to accumulate energy in the physical activity are stress hormones including thyroid hormones or enzymes expressed by genes as ACE. (Heshmat et al., 2010), Burger (2004)

Inherited defects of thyroid hormone metabolism has been discussed by Dumistrescu and Refetoff (2011) They came to the conclusion that the only known inherited defect of intracellular thyroid hormone metabolism is caused by mutations in the SBP_2 gene affecting selenoprotein synthesis, among which the typical laboratory findings are high T_4 , Low T_3 , high $r T_3$, normal or slightly elevated serum TSH, decreased serum se and decreased selenoprotein levels and activity in serum and tissues. The clinical phenotype is complex; affected individuals may have delayed growth and puberty, and in severe cases failure to thrive, mental retardation, infertility and myopathy. SBP_2 defects could have as yet undetermined consequences and the identification of additional patients, and their long term follow up are important in further characterizing this recently described defect. So, we investigated the role of endurance exercise program, in hyperthyroidism in relation to ACE genotype.

Methods

The study included 10 female thyroid patients (hyperthyroid) and 10 female control group. Their baseline characteristic are reported in

Table (1)

Variables	Thyroid patients (n=10)	Control (n=10)
Age, Y	35 \pm 4	37 \pm 5
Gender	Female	Female
Height, cm	162 \pm 7	164 \pm 9
BMI	23.4 \pm 5.6	22.3 \pm 4.5
Weight, Kg	63 \pm 11	61 \pm 12
Blood pressure mm/Hg	150/90	122/78
Pulse Rate count/min	88 \pm 6	78 \pm 7

Values M \pm SD significant $P < 0.05$

Venous blood was obtained from all participants for genomic DNA and ACE genotype, pulse rate, B/P, BMI. Thyroid function test (TSH,

$F.T_{3,4}$, T_{T3} $T_{4,}$) each participant underwent a medical history and Physical examination. The presence of Thyroid dysfunction was determined by lab test and clinical assessment.

Identification of ACE genotype

Genomic DNA was extracted, then DNA fragments were amplified by PCR, genotype was performed by Quiaxcel. Including concentration and pulse rate, BP in each case of participants, diseased and control.

The main symptoms of hyperthyroidism in the examined patients of the study :

- a high state of excitability.
- intolerance to heat.
- increased sweating.
- weight loss.
- varying degrees of diarrhea.
- muscle weakness.
- extreme fatigue.
- nervousness.
- inability to sleep.
- tremors of the hands.
- some patient develop some degree of protrusion of the eyeballs (exophthalmos).
- some other patients with increased thyroid gland (Barret et al, 2010).

The PCR conditions were determined using a primer pair of sense 5-CTG GAG ACC ACT CCC ATC CTT TCT – 3 and antisense 5-GAT GTG GCC ATC TTC GTC AGA- 3. PCR amplification was carried out in a total volume of 25 ul containing 50ng template DNA, Hormone estimation was carried out using Elisa technique.

Endurance training program after Grandys (2008).

Statistical analysis: Genotypes among the patients and controls were obtained by direct enumeration based on PCR results.

3. Results

The results of hormones were reported as Means \pm SD.

T test was also performed between two groups, significant results reported $P < 0.05$.

Table (2) comparison of the studied parameters in control and hyperthyroid groups.

Parameters	TSH m iu/L	F T_4 (pmol/L)	F T_3 (pmol/L)	T T_4 (mmol/L)	T. T_3 (mmol/L)
Groups					
control n = 10	2.4 \pm 0.23	16.4 \pm 4.1	1.80 \pm 0.5	81 \pm 11.2	2.6 \pm 0.2
hyperthyroid patients n = 10	(*) 0.1 \pm 0.02	(*) 32.1 \pm 6.2	(*) 7.9 \pm 2.1	(*) 90 \pm 13.0	(*) 2.9 \pm 0.3

Values given as the mean + SD * Significant difference at $P < 0.05$

Table (3) comparison of the studied parameters in hyperthyroid group before and after endurance exercise program.

Group \ Parameter	TSH miu/L	F T ₄ (pmol/L)	F T ₃ (pmol/L)	T T _u (mmol/L)	T. T ₃ (mmol/L)
hyperthyroid before endurance exercise	0.1±0.02	32.1±6.2	7.90±2.1	90±13.0	2.9±0.3
hyperthyroid after endurance exercise	(*) 1.3±0.6	(*) 25.2±4.4	(*) 2.4±0.7	84±6.4	2.8±0.4

Values M ± SD Significant at P < 0.05

Table (4) Comparison of the pulse rate and blood pressure in hyperthyroid group before and after endurance exercise.

Group \ Parameter	Pulse rate Count/min	Blood Pressure mm Hg
hyperthyroid before exercise	88 ± 6	150/90
hyperthyroid after exercise	82 ± 5 *	138/84 *

Values M ± SD; significant P < 0.05




Table (2) revealed a decreased TSH, and increased F.T₄, T₃, T₄, T₃ in hyperthyroid subjects.

- The results indicated (table3) a decrease in T_{T3}, and T_{T4}, hormones after the endurance training program.
- TSH increased after endurance exercise in hyperthyroid cases.
- The results also revealed a decreased T_{T3}, T_{T4}, after endurance physical effort.

As for ACE genotype, the results indicated that the ACEDD and ACEID represents a major association with abnormalities of thyroid function and may have a pathogenic role in thyroid glands and needs to be addressed in future studies. In control group the ACE genotype recorded revealed ACEDD (25%) I/D (50%) ACEII (25%)

4. Discussion

TSH acts on thyroid metabolism through its stimulation of thyroid hormones at all stages such as iodine uptake, organization and coupling, it also enhances the release of stored thyroid hormones, and increases DNA content, RNA and translation of proteins together with stimulation of glycolysis and activates adipose tissue lipase to enhance the release of fatty acids (Chatlerjea and Rana Shinde, 2006) Yen (2001).

The data presented in table (2) indicated a low TSH concentration (0.1 mu/ L.) this result is in agreement with that of Smith et al (1990)

Mona Abou Zahra et al (2006) and Stockigt (1996). They reported that the measurement of TSH in a basal blood sample provides the single most sensitive, specific and reliable test of thyroid status. In case of hyperthyroidism TSH is below 0.1 m u/L. and a normal TSH excluded a primary thyroid disorder (Cooper, 2003).

As for F.T₃, T₄ and T.T₃ T₄, data presented in Table (2) indicated a higher level of both F.T₃ and T₄ and T.T₃, T.T₄. The measurement of thyroid hormones free or total provide the most reliable indication of thyroid status, as free and total T₃, T₄ can help assess the severity of thyroid disease and distinguish subclinical from overt disease. Smith et al (1990) reported that free thyroid hormone measurements correlate more closely with thyroid status than total hormone measurements, which are heavily influenced by changes in the concentration of thyroid hormone – binding proteins. Murray et al., (2009) stated that one half to two thirds of T₃ and T₄ in the body is in an extrathyroidal reservoir most of this circulates in bound form, bound to a specific binding protein, Thyroxine – binding globulin (TBG), is a glycoprotein with a molecular mass of 50 KDa, binds T₄ and T₃ and has the capacity to bind 20 ug/ed of plasma. Under normal circumstances, TBG, binds nearly all of the T₄ and T₃ plasma, and it binds T₄ with greater affinity than T₃. Thus, in spite of the great differences in total amount, the free fraction of T₃ approximates that of T₄ and given that T₃ is intrinsically more active than T₄, most biologic activity is attributed to T₃, T BG does not bind to other hormones. (Guyton and Hall, 2006 Dayan 2001)

From the conversion of Angiotensin I into angiotensin II by ACE. La Churie et al. (1995) stated that contracting with the lack of relation between ACE gene polymorphism and blood pressure level, a large case – control study has shown that the deletion marker allele of the ACE gene was associated with an increased risk of myocardial infarction. While, Ohishi, (1993) identified that the D allele has been associated with growth of vascular smooth muscle at the site of coronary angioplasty and human cardiac hypertrophy in response to exercise (Montgomery et al., 1997). Kumar et al., (2012) investigated seven polymorphisms including ACE gene and essential hypertension and found a possible contribution of ACE and Nos in the complex etiology of hypertension. As for ACE genotype, the results indicated 80% ACEDD, 20% in case of ACEID with a higher Blood pressure in case of ACEDD compared to ACEID genotype. In case of the control group they have normal blood pressure and ACEDD 25%,

ACEID 50%, ACEII 25% This was also reported by Jalil et al (2004) . Heshmat et al., (2010).

Andreas (1997) stated that ACE is a key component of the renin angiotensin system thought to be important in the pathogenesis of hypertension and other cardiovascular disease, Tiret et al., (1992) recorded that the D allele of ACE genes is characterized by higher ACE in the blood. Hashimoto et al (2001) reported that the increase in blood pressure noticed in subjects having ACEDD might results. After an exercise training program of low intensity endurance exercise for 3 months, there was an elevated TSH concentration together with decreased level of F.T₃, T₄ and T. T₃, T₄ (Table 3) .

The exercise training program induced a positive effect on thyroid hormone parameters in hyperthyroid patients, the positive effect of endurance exercise might be due to decreased stresses on the patients and hence a relaxation of muscles of blood vessels which induced a higher stimulation on pituitary and thyroid glands, with the end results a positive effect and higher metabolic action. Fricker et al. (2003) and Wiesli et al (2003) reported that thyroid dysfunction is associated with changes in cystatin C which can be used as marker for thyroid patients due to direct thyroid hormones upon it.

The decreased thyroid hormones and elevated TSH concentration was noted after exercise in many studies such Malik (2011), Maha et al (2011), Mohamed et al (2009). Grandys et al (2008), Figen et al (2005), Loucks et al (1993). They stated that the decreased F.T₃ might be caused due to R T₃ formation which act as a brake if the body needs lesser energy, which direct the thyroid gland to synthesize more reverse T₃ (R T₃) than T₄ .

Table (4) revealed that endurance exercise training induced a significant decrease in pulse counts and blood pressure, which indicated a better fitness state to the hyperthyroid patients. It was also noted that there was an association with the occurrence of hypertension and hyperthyroid patients and the prevalence of ACE genotype DD and ID. This was in accordance with Wu et al (2011) and Siklar et al (2011).

Conclusion

- 1- ACEDD, ID genotypes predominate hyperthyroidism and may have a pathogenic role in thyroid gland.
- 2- TSH decrease is the main lab test of hyperthyroidism.
- 3- Endurance training might have a positive effect alone or together with antithyroid drugs in treating hyperthyroidism

Recommendation

To use ACE genotype to evaluate hyperthyroidism, and thyroid hormones as indicators of training efficiency.

References

- 1- Andreas B. Hans, K, Friedrich, L (1997) ACE and ANG gene polymorphisms, plasma levels, Cardiac dimensions Hypertension 29: 165 – 170.
- 2- Barrett, K., Barman, S., Boitana, S. (2010) Ganong Medical Physiology MC graw Hill Lange USA.
- 3- Bungler, A (2004) Environment and thyroid function J Clin. Endocrinol Metab 89 : 1526.
- 4- Chatterjee, M. and Rana Shinde (2006) Medical Biochemistry, Jaypee, India.
- 5- Cooper D. (2003) Hyperthyroidism Lancet 362: 459.
- 6- Dayan. C. (2001) Interpretation of thyroid function tests Lancet 357: 619.
- 7- Dumitrescu , A. and Refetob, S. (2011) Inherited defects of thyroid hormone metabolism Ann Endocri., 72:95-98.
- 8- El Taib, S. (2012) the concentration level of some thyroid and adrenal gland hormones in athlete blood plasma PHD Dissertation, Alexandria Univ. Egypt.
- 9- El Shafei, M., Farres, M. Shahin, R. (2011) Evaluation of angiotensin converting enzyme gene polymorphism and susceptibility to bronchial asthma among Egyptians Aller Immunopathol 1:25.
- 10- Figen, C. (2005) Exercise intensity and its effect on thyroid hormones Neuroend. Letters 26: 68.
- 11- Fricker, M. Wiensli, P. Brandelle, M. (2003) Impact of thyroid dysfunction on serum cystatin C Kid. Int. 63, 1944- 1947.
- 12- Grandys, J., (2008) the effect of endurance training on muscle strength in young, healthy men in relation to hormonal status J. of physiol. And pharmacology 59, 89 – 103.
- 13- Guyton, A. and Hall, J. Medical physiology El Sevier, Saunders, USA.
- 14- Hashimoto, Y. Futamara, A., Nakara. H. (2001) Relationship between response of glutamyl transpeptidase to alcohol drinking and risk factors for coronary heart disease Athero Sclerosis 138: 465- 670.
- 15- Heshmat, H, Azab, A. and El Tantawy, P (2010) ACE and ACE gene polymorphism and some cardiac parameters of fencing, basket ball, endurance players World J. of Sport Sc. 3: 100- 106.
- 16- Jalil, J. Ocaranz M. Oliveri, (2004) Neutral endopeptidase and ACEI gene polymorphism in Human J. hypertension 18: 119-125.
- 17- Joshi, M., Wattanapitay Akal, S. Bauer, J. (2011) effect of endothelial gene polymorphisms on cellular responses, to hyper glycaemia : Role of ACE (I/D) Dib Vasc Dis Res 8: 276 – 283.
- 18- Kumar, V., Selby, A. Rankim, D. (2012) Age Related differences in dose response of muscle protein synthesis to resistance exercise in young and old men J. physiol. 582: 814-821.
- 19- La Churie, M., Aziz, I., Menand, J (1995) ACE gene polymorphism has no influence on the circulating Renin Angiotensin – Aldosterone System of blood pressuring normotensive subjects Circulation 15: 91: 2933-2942.
- 20- Louck S., A. (1993) introduction and prevention of low T3 Syndrome The Am. Physiol. Soc., 23: 112.
- 21- Maha, A. (2011) Estimation of post exercise thyroid hormones in trained Student of physical education college underwent submaximal exercise Tikrit J. of Pharmaceutical Sciences 1:202.
- 22- Malik, B. (2011) The effect of maximal aerobic exercise on cortisol and thyroid hormones in male hockey Players. African J. Pharmacy and pharmacology 5: 2002- 2006.
- 23- Mohamed, J. (2009) The alteration of serum thyroid hormone and its stimulating in nano scale athletics men Univ. of Kashan, Iran.
- 24- Mona Abou Zahra. Walid, Z. and Omayma, H. (2005) Impact of thyroid dysfunction clearance Egypt J. of Biol. and Mol. Biol. 24:13.
- 25- Montgomery H., Vlarckson, P. and Dollery, C. (1997) Association of angiotensin Converting enzyme gene (I/D) in response to physical training Circulation 96: 741- 747.
- 26- Murray, R., Bender, D. Weil, A. (2009) Harper illustrated biochemistry Lange, New York.
- 27- Ohishi, H. (1993) ACE I/D angioplasty in response to exercise Hypertension 12: 87-92.
- 28- Pavlic, D., Ristic, Flego, V. (2012) ACE I/D gene polymorphism lung – cancer patients gene test Mol. Biomarkers 12:115.
- 29- Rigat, B., Hulent, C. Corvol, P. (1992) PCR detection of the I/D polymorphism of the human ACE gene Nucleic ACIDS Res. 20:1433.
- 30- Siklar, Z., Berberoglu, M. Savan, S. (2011) Contribution of clinical, metabolic and genetic factors on hypertension in obese children and adolescents J Pediatr. Endocrinol Metab 24:21-24.
- 31- Smith, A., Bekett, G., Walker, S. (1990) Lecture notes on clinical Biochemistry 6th ed. Blackwell SC. Publ., London.
- 32- Stockigt, J. (1996) Guidelines on diagnosis and monitoring of thyroid disease Clin Chem. 42:188- 192.
- 33- Tired, L., Rigat, B., Soubrier, F. (1992) Evidence from combined Segregation and linkage analysis a variant of ACE gene control plasma ACE Levels Am. J. Genet. 51:197-205.
- 34- Wienli, P., Schueglr. B., Spina, G. (2003) Serum cystatin C is sensitive to small change in thyroid function Clin chem. Acta 338:87 – 90.
- 35- Wa, F., Li, G. Songx (2011) Relationship between ACE gene polymorphism and essential hypertension Zhongguo Dang Dai ER 13: 883 – 885.
- 36- Yen, P., (2001) Physiological and Molecular basis of thyroid hormone action. Physiol Rev 81: 1097.

1/15/2013