

Molecular Characterisation of Lipolytic Fungi Isolated From Meat Products

Ashraf A. Abd El-Tawab¹; Eman M. El-Diasty²; Gamal R. Hassab El-Naby²; Fatma I. El-Hofy¹; Abd El-hady M. Fkhreldeen³

¹Bacteriology, Immunology and Mycology Department, Faculty of Veterinary Medicine, Faculty of Veterinary Medicine, Benha University, Egypt

² Mycology Department, Animal Health Research Institute Dokki, Giza, Egypt

³ M.V.Sc. Cairo University, Egypt

dr_noor_106@yahoo.com

Abstract: This work was carried out to evaluate the fungal quality of luncheon, basterma and minced meat sold in different groceries and supermarkets in Gharbia Governorate under different trade names, as well as to evaluate lipolytic activity of isolated fungi and molecular characterization of some of this lipolytic fungi isolated from meat products. The mean total mould counts were $3 \times 10^2 \pm 1.2 \times 10^2$, $2.7 \times 10^2 \pm 3.9 \times 10$ and $5.2 \times 10^3 \pm 1.9 \times 10^3$ cfu/g for luncheon, basterma and minced meat, respectively. In the examined samples, 7 mould genera were identified. The isolated mould species were *Aspergillus* sp, *Alternaria* sp, *Endomyces* sp, *Eurotium* sp, *Penicillium* sp, *Monascus* sp and *Geotrichum* sp were isolated from the examined samples at varying percentages. *Aspergillus* was represented by 4 species, *A. flavus* and *A. niger* were isolated from examined meat product, while *A. ochraceus* and *A. roseoglobulus* were isolated from basterma only. A total of 55 isolates belonging to 17 species were tested for their ability to produce lipolytic enzyme. Four lipolytic *A. niger* isolates were subjected to PCR identification, and were sequenced in both directions. Sequences were analysed and aligned by Clustal method using the program of DNA star (Laser-gene, Wisconsin, USA).

[Ashraf A. Abd El-Tawab; Eman M. El-Diasty; Gamal R. Hassab El-Naby; Fatma I. El-Hofy; Abd El-hady M. Fkhreldeen. **Molecular Characterisation of Lipolytic Fungi Isolated From Meat Products.** *J Am Sci* 2019;15(4):8-16]. ISSN 1545-1003 (print); ISSN 2375-7264 (online). <http://www.jofamericanscience.org>. 2. doi:[10.7537/marsjas150419.02](https://doi.org/10.7537/marsjas150419.02).

Key word: *A. niger*, lipolytic fungi, meat product

1. Introduction

Undoubtedly the main source of animal protein is meat and its products and it is also a high source of several vitamins and minerals, and so meat products are considered a favourable food as it's easy to buy, fast to cook so it's the first choice for many people to eat. Meat products may contaminate with one of the most dangerous microbial hazards represented in moulds. (Morshdy et al., 2015)

Meat products become contaminated during processing and handling operations. Dijksterhuis and Samson. (2002)

Lipases (triacylglycerol acylhydrolase, EC 3. 1. 1. 3) are hydrolytic enzymes that catalyze the hydrolysis of fats and oils to glycerol and free fatty acids at the oil-water interface (Jooyandeh et al., 2009). A considerable number of fungal and bacterial lipases have been commercially produced with the former being preferable, because fungi generally produce extracellular enzymes, Extracellular secretion of lipase enzyme has been well studied for a number of fungi, primarily Zygomycetes, hyphomycetes, and yeasts. In meat, lipid hydrolysis can take place enzymatically or non-enzymatically. The enzymatic hydrolysis of fats is termed lipolysis or fat deterioration and is governed by specific enzymes

such as lipases, esterases and phospholipase. Lipolytic enzymes could either be endogenous of the food product (such as milk) or derived from psychrotrophic microorganisms (Ghaly et al., 2010). During lipolysis, lipases split the glycerides forming free fatty acids which are responsible for common off-flavour, frequently referred to as rancidity (Huis in't Veld, 1996 and FAO, 1986). The main enzymes involved in meat lipid hydrolysis are phospholipase A1 and phospholipase A2 (Toldra, 2006). Lipid hydrolysis process is regiospecific and involved three steps of biosynthetic pathway: cleavage of triacylglycerol, acyl migration and cleavage of 1-monoacyl-sn-glycerol. (Belitz et al., 2009; Christie, 2010)

Internal Transcribed Spacer (ITS) regions have been successfully used to generate specific primers capable of differentiating closely related fungal species. Analysis of the ITS is also sensitive enough to identify intraspecific variation that is not apparent from morphological analysis. (Abd-El salam et al., 2004)

This study was carried out to throw light on the incidence of lipolytic fungi in meat products with molecular characterisation of this lipolytic fungi existing in such products.

2. Material and methods

Collection of samples

A grand total of 60 random samples representing table luncheon (20), basterma (20) and minced meat (20) were collected under different trade names then were kept in sterile polyethylene bags and preserved in an ice box then transferred to the laboratory under complete aseptic condition without undue delay to be examined mycologically.

Food homogenate preparation: according to APHA (2001).

Determination of total mould count according to the technique recommended by (ISO, 2008):

Identification of mould isolates: according to the technique recommended by Pitt and Hocking (2009), Frisvad and Samson. (2004)

Detection of lipolytic activity: according to Sunitha et al., (2013)

We used the sensitive plate assay for detection of lipase activity in growing cultures (10, 11). Detection of lipase activity was performed on Peptone Agar medium supplemented with Tween 20 separately sterilized and added 1% to the medium. At the end of the incubation period, a visible precipitate around the colony due to the formation of calcium salts of the lauric acid liberated by the enzyme indicated positive lipase activity.

Fungal Mycelium Preparation and DNA Isolation: Fungal mycelium was prepared from pure culture using 50 ml of SDA broth in 100 ml conical flasks and was incubated at $25 \pm 1^\circ\text{C}$ for 7 days. Mycelia from 50 ml broth were harvested by filtration through Whatman sterile filter paper and the fungal genomic DNA was isolated (Rohlf, 1998). After DNA extraction, lipase-producing fungi were identified via ITS-PCR of rDNA region with ITS1 and ITS4 primers. PCR amplification conditions for *A. niger* were: 5 min initial step followed by 38 cycles at 94°C for 1 min, 50°C for 1 min and 72°C for 1.5 min and a final extension step at 72°C for 5 min. Amplification products were electrophoresed in agarose gels (3% w/v) (Agarose, Sigma, USA). And stained with

ethidium bromide using Gene Ruler 100bp DNA Ladder (Fermentas Company, Cat.No.SM0243, USA). Finally, we made sequencing to the PCR product on GATC Company by use ABI 3730xl DNA sequencer by using forward and reverse primers. Only by combining the traditional Sanger technology with the new 454 technology, can genomes now be sequenced and analyzed in half the usual project time, with a considerable reduction in the number of coatings and gaps. In addition, considerable cost advantages now make genome sequencing with the 454-technology accessible to the research community. Sequences were analyzed and aligned by Clustal method using the program DNA star. (Laser-gene, Wisconsin, USA)

3. Results and Discussion

Meat is preferred by millions of people all over the world as a major source of animal protein. But meat is considered a favourable media for growth of various pathogens, so could be spoiled rapidly. The decomposition of different constituents of meat including protein and fat resulting in bad odours and other unfavourable changes. Therefore, spoilage of meat and meat products should be limited and controlled to protect the human consumers from infection and allergy (Aksoy et al., 2014). In the present study, the mean mould count was $3 \times 10^2 \pm 1.2 \times 10^2$, $2.7 \times 10^2 \pm 3.9 \times 10^1$ and $5.2 \times 10^3 \pm 1.9 \times 10^3$ cfu/g in luncheon, basterma and minced meat, respectively as described in Table (1). The results of luncheon samples nearly similar to those obtained by Shaltout and Salem (2000) and Elsayed et al (2018) reported that the mean mould count in luncheon and minced meat was $1.3 \times 10^2 \pm 2.1 \times 10^1$ cfu/g and $2.8 \times 10^2 \pm 4.3 \times 10^1$ cfu/g, respectively. while higher prevalence was recorded by El-Tabiy (2006) who reported that the mean mould count in luncheon and minced meat were $1.3 \times 10^4 \pm 9.3 \times 10^3$ and $2.1 \times 10^6 \pm 6.1 \times 10^5$ cfu/g, respectively. Hussein (2008) who reported that mean mould count of sausage. ($2.26 \times 10^2 \pm 0.58 \times 10^2$ cfu/g)

Table (1): Incidence of mould isolated from meat products (No=20 for each product):

Meat products	No of examined samples	No of positive mould samples	Min.	Max.	Mean \pm SE
Luncheon	20	9	3×10^1	8.5×10^2	$3 \times 10^2 \pm 1.2 \times 10^2$
Basterma	20	20	2×10^1	5.2×10^2	$2.7 \times 10^2 \pm 3.9 \times 10^1$
Minced meat	20	14	3.3×10^2	7.8×10^3	$5.2 \times 10^3 \pm 1.9 \times 10^3$

Mould contamination of meat and meat products may occur during slaughtering of the animals, transportation, or during processing of meat products through the use of contaminated equipments or contaminated additives and spices which considered the most important source of mould contamination in meat products. (Jay et al., 2005)

The data obtained from table (2) declared that 7 mould genera could be isolated and identified from the examined meat samples. The identified mould genera were; *Aspergillus*, *Alternaria*, *Endomyces* spp, *Penicillium* spp, *Monascus* spp *Geotrichum* spp. In addition, that 4 *Aspergillus* species could be identified from meat product samples. The most predominant

species were; *Aspergillus flavus* and *Aspergillus niger* which were present in 4 (20%), 4 (20%), 2(10 %), 2 (10 %), 4 (20%) and 3(15%), respectively, in Luncheon, Basterma and Minced meat. *A. flavus* and *A. niger* were present in all examined meat samples. *A. ochraceus* and *A. roseoglobulus* were only present in Basterma. Incidence of *Penicillium* species isolated from examined meat samples revealed that *P. citreonigrum*, *P. glabrum*, *P. polonicum*, *P. citrinum* and *P. simplicissim* were the most predominant species isolated from Luncheon with frequency of 1(5%). In addition, the result achieved in table (2) it is obvious that *P. crusteoum*, *P. paneum* and *P. brevicompactum* were isolated from the Basterma samples only.

The results illustrated in table (3) showed *A. niger*, *A. flavus*, *A. ochraceus*, *P. citreonigrum*, *P. glabrum*, *P. cirtinun*, *P. crusteoum*, *P. paneum*, *P. brevicompactum* and *P. simplicissim* were having lipolytic activity with a visible precipitate around the colony due to the formation of calcium salts of the lauric acid liberated by the enzyme indicated positive lipase activity. These results nearly similar to those obtained by **Nasser (2002)**, **Shaltout (2002)**, **Subash, et al. (2005)**, **Soliman and Shalaby (2008)**, **Korashy and Wahbba (2008)**, **El-Diasty and Salem (2009)**, **Ouf et al. (2010)** and **Shaltout et al. (2014)**

Table (2): Incidence of isolated mould from meat products (No=20 for each product):

Isolated mould genera	Luncheon		Basterma		Minced meat	
	No	%	No	%	No	%
<i>Aspergillus</i> spp	6	30	14	70	5	25
<i>A. flavus</i>	4	20	4	20	2	20
<i>A. niger</i>	2	10	4	20	3	15
<i>A. ochraceus</i>	-	-	5	25	-	-
<i>A.roseoglobulus</i>	-	-	1	5	-	-
<i>Alternaria</i> spp	1	5	2	10	-	-
<i>Endomyces</i> spp	-	-	-	-	7	35
<i>Eurotium</i> spp	2	10	-	-	-	-
<i>Penicillium</i> spp	5	25	6	30	-	-
<i>P. citreonigrum</i>	1	5	-	-	-	-
<i>P. glabrum</i>	1	5	-	-	-	-
<i>P. polonicum</i>	1	5	-	-	-	-
<i>P. citrinum</i>	1	5	-	-	-	-
<i>P. crusteoum</i>	-	-	2	10	-	-
<i>P. paneum</i>	-	-	2	10	-	-
<i>P. brevicompactum</i>	-	-	2	10	-	-
<i>P. simplicissim</i>	1	5	-	-	-	-
<i>Monascas</i> spp	1	5	1	5	-	-
<i>Geotrichum</i> spp	1	5	-	-	4	20

These PCR systems described to date are useful only in identifying the genus *Aspergillus* as a whole or the single species. The ITS region contains variable elements that allow for sequence-based identification of *Aspergillus* species (**Iwen et al., 2002**); therefore, the region offers a possible template for design of species-specific primers for identification of the major pathogenic species. Most previous publications on PCR-based detection or identification systems for *Aspergillus* spp. were based on using 18S or 28S rDNA as target DNA. However, the sequences in these regions are conserved across a wide range of fungi; it is therefore difficult to design truly species-specific primers. As reported previously, the more variable ITS regions have proven more useful for identification of fungal species **Makimura. (2001)**

Polymerase Chain Reaction (PCR) amplification of ITS region of rRNA genes with ITS1 and ITS 4 primers yielded distinct DNA bands for all representative isolates investigated DNA sequences of *Aspergillus niger* identified by similarity searches in the GenBank gave.

Molecular biological identification systems for pathogenic aspergilli have been suggested as a solution to this problem: for example, a PCR based diagnostic method for detecting the genus *Aspergillus* using 18S rDNA has been designed (**Makimura et al., 1994 and Yamakami et al., 1996**). Systems have also been described for specific detection of *Aspergillus niger* with primers based on regions of the 28S rDNA (**Henry et al., 2000 and Zhao et al., 2001**) or of the internal transcribed spacer (ITS) 1 and 2 regions of ribosomal DNA (rDNA). These PCR systems

described to date are useful only in identifying the genus *Aspergillus* as a whole or the single species *A. niger*. The ITS region contains variable elements that

allow for sequence-based identification of *Aspergillus* species.

Table (3): Lipolytic activity of mould on tween 80 media:

strains	No of examined strains	positive strains		No of negative strains
		No of positive strains	Diameter of inhibition zone (mm)	
<i>A.niger</i>	15	15	16-17	-
<i>A.flavus</i>	16	9	10	7
<i>A.ochraceus</i>	5	5	19	-
<i>A.roseoglobulus</i>	1	-	-	1
<i>P.citreonigresum</i>	4	4	9	-
<i>P.glabrum</i>	2	2	7	-
<i>P.polonicum</i>	1	-	-	1
<i>P.cirtinun</i>	1	1	12	-
<i>P.crusteoum</i>	2	2	17	-
<i>P.paneum</i>	3	3	12-13	-
<i>P.brevicompectum</i>	2	2	10-11	-
<i>P.simplicicissim</i>	3	3	7	-
<i>Alternaria sp</i>	3	-	-	3
<i>Endomyces sp</i>	7	-	-	7
<i>Euroteium spp</i>	2	2	6	-
<i>Monoscasp spp</i>	2	-	-	2
<i>Geotrichum spp</i>	5	5	13-14	-




Figure (1): 1.5 % of agarose gel electrophoresis of PCR product with ITS for *A.niger*

Lane1 Molecular weight marker, Lane2-4 *A. niger*.

Three *A.niger* isolates has lipolytic activity on peptone agar medium supplemented with tween 20 were examined by molecular methods polymerase chain reaction (PCR) with using ITS primer. PCR products of *A.niger* strains were positive on agarose

gel electrophoresis of PCR amplification products. (Figure 1)

The PCR products of the one *A.niger* isolate was sequenced and aligned with references in the NCBI database. The sequences had 98.1% identity with *Aspergillus* spp. sequences deposited in the NCBI database. Nucleotide analysis in figure (2) showed greater than 98% similarity for the entire sequence of the one sequenced strain *A.niger*-AAE-EG017 in comparison of 10 strains of *Aspergillus* Spp. The sequence similarity for the strains of *A.niger*-SAM6, *A.niger*-36, *A.tubingensis*-ND9, *A.niger*-129B, *A.niger*-FIS17, *A.tubingensis*-G4-14, *A.tubingensis*-F43-01, *A.niger*-AL-29, *A.awamori*-ND6 and *A.niger*-Egypt-BSU-5 were 100 %, 100%, 100%, 100%, 99.7%, 99.4 %, 99.7%, 99.7% and 99.7 % . (figure 2).

Phylogenetic analysis showed that *A.niger*-AAE-EG017 accession no. (MH590624) was isolated but in different branches, away from *A.niger*-SAM6, *A.niger*-AL-29, *A.niger*-36, *A.niger*-129B, *A.niger*-FIS1 and *A.niger*-Egypt-BSU-5 but remained in the same cluster. While *A.niger*-AAE-EG017 accession no. (MH590624) still in the same group of *A.tubingensis*-G4-14. (figure 3)

Of the strain of *A. niger* studied, ITS could be amplified from three strains. The *A.niger* strain is readily distinguished from those of the other species by the 17 base insertion (TCCAGCATCGAATGGG) at nucleotides 1–17, and their distinctive nucleotide changes (AGGA) at the following positions: 4-15-16-17. Some of these variations result in nucleotide lead to amino acid changes. (Figure 4)

		Percent Identity												
		1	2	3	4	5	6	7	8	9	10	11		
Divergence	1	■	98.1	98.1	98.1	98.1	98.1	98.3	98.3	98.1	97.8	98.0	1	A.nig-AAE-EG017
	2	2.0	■	100.0	100.0	100.0	100.0	99.7	99.7	100.0	99.7	100.0	2	A.niger-SAM6
	3	2.0	0.0	■	100.0	100.0	100.0	99.7	99.7	100.0	99.7	100.0	3	A.niger-36
	4	2.0	0.0	0.0	■	100.0	100.0	99.7	99.7	100.0	99.7	100.0	4	A.tubingensis-ND9
	5	2.0	0.0	0.0	0.0	■	100.0	99.7	99.7	100.0	99.7	100.0	5	A.niger-129B
	6	2.0	0.0	0.0	0.0	0.0	■	99.7	99.7	100.0	99.7	100.0	6	A.niger-FIS17
	7	1.7	0.3	0.3	0.3	0.3	0.3	■	99.4	99.7	99.4	99.7	7	A.tubingensis-G4-14
	8	1.4	0.0	0.0	0.0	0.0	0.0	0.3	■	99.7	99.4	99.7	8	A.tubingensis-F43-01
	9	2.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	■	99.7	100.0	9	A.niger-AL-29
	10	2.3	0.3	0.3	0.3	0.3	0.3	0.6	0.3	0.3	■	99.7	10	A.awamori-ND6
	11	2.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.3	■	11	A.niger-Egypt-BSU-5
	1	2	3	4	5	6	7	8	9	10	11			

Figure (2): Nucleotides identity of sequences of different *Aspergillus* spp.




Figure (3): Phylogenetic tree of **Nucleotides** sequences of different *Aspergillus* spp.

Majority	GAAAGGCAGCGGGCCACCGCGTCCGATCCTCGAGCGTATGGGGCTTTGT	
	-----+-----+-----+-----+-----+-----	
	210 220 230 240 250	
	-----+-----+-----+-----+-----+-----	
A.nig-AAE-EG017	GAAAGGCAGCGGGCCACCGCGTCCGATCCTCGAGCGTATGGGGCTTTGT	250
A.niger-SAM6	242
A.niger-36	242
A.tubingensis-ND9	242
A.niger-129B	242
A.niger-FIS17	242
A.tubingensis-G4-14	242
A.tubingensis-F43-01	241
A.niger-AL-29	242
A.awamori-ND6	242
A.niger-Egypt-BSU-5	242
Majority	CACATGCTCTGTAGGATTGGCCGGCGCCTGCCGACGTTTCCAACCATT	
	-----+-----+-----+-----+-----+-----	
	260 270 280 290 300	
	-----+-----+-----+-----+-----+-----	
A.nig-AAE-EG017	CACATGCTCTGTAGGATTGGCCGGCGCCTGCCGACGTTTCCAACCATT	300
A.niger-SAM6	292
A.niger-36	292
A.tubingensis-ND9	292
A.niger-129B	292
A.niger-FIS17	292
A.tubingensis-G4-14	292
A.tubingensis-F43-01	291
A.niger-AL-29	292
A.awamori-ND6C	292
A.niger-Egypt-BSU-5	292
Majority	TTTCCAGGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACCTAAGCAT	
	-----+-----+-----+-----+-----+-----	
	310 320 330 340 350	
	-----+-----+-----+-----+-----+-----	
A.nig-AAE-EG017	TTTCCAGGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACCTAAGCAT	350
A.niger-SAM6	342
A.niger-36	342
A.tubingensis-ND9	342
A.niger-129B	342
A.niger-FIS17	342
A.tubingensis-G4-14	342
A.tubingensis-F43-01	341
A.niger-AL-29	342
A.awamori-ND6	342
A.niger-Egypt-BSU-5	342
Majority	ATCAATAAGGCGGAGGAAXXX	
	-----+-----+-----	
	360 370	
	-----+-----+-----	
A.nig-AAE-EG017	ATCAATAAGGCGGAGGAAAAA	371
A.niger-SAM6	361
A.niger-36	360
A.tubingensis-ND9	360
A.niger-129B	360
A.niger-FIS17-	362
A.tubingensis-G4-14-	359
A.tubingensis-F43-01-	360
A.niger-AL-29-	359
A.awamori-ND6	360
A.niger-Egypt-BSU-5	356

Conclusion

This investigation has shown that all meat products are important contributors to the transmission of *Aspergillus* spp (*A. niger*) which had lipolytic activity and extract ochratoxin A which cause economic loss and toxicity.

References

1. Abd-Elsalam, K.A.; Aly, I.N.; Abdel-Satar, M.A.; Khalil, M.S. and Verreet, J.A. (2004): PCR identification of *Fusarium* genus based on nuclear ribosomal DNA sequence data. *Afr. J. Biotechnol.*;2(4):82-5.
2. Aksoy, A.; Sezer, C.; Aydin, D.B. and Güven, A. (2014): The effect of some natural antimicrobial substances on the shelf life of beef. *Israel J. of Vet. Med.* 69(3):114-122.
3. APHA (2001): American Public Health Association: Compendium of methods for the microbiological examination of foods. 4th Ed. Eds. Downes, F.P. and K. Ito. Sheridan Books Inc., Washington D.C., USA.
4. Belitz, H.D.; Grosch, W. and Schieberle, P. (2009): *Food Chemistry* (4th Edition). Springer-Verlag Berlin Heidelberg, Germany, pp: 596.
5. Christie, W.W. (2010): Triacylglycerols Part 2. Biosynthesis and Metabolism. AOCs Library, <http://lipidlibrary.aocs.org/Lipids/tag2/index.htm>
6. Dijksterhuis, J. and Samson, R.A. (2002). Food and crop spoilage on storage. In F. Kempken (Ed.), *the Mycota. Vol XI Agricultural Applications*. Springer-Verlag Berlin Heidelberg, Germany, pp 39-52.
7. El-Diasty, E. M. and Salem, R.M. (2009): Incidence of lipolytic and proteolytic fungi in some milk products and their public health significance. *Arab J. Biotech.*, Vol. 12, No. (1) Jan. (2009): 49-56.
8. Elsayed, M. E.; Abdelazeem M. A.; El-Diasty, E. M.; Abouelmaatti, R. R. and Abbas, S. M. (2018): Prevalence of *Aspergillus* spp and Aflatoxins in luncheon, minced meat and sausage. *Global Animal Science Journal-GASJ* 6(2):17-23.
9. El-Tabiy, A. A. (2006): Mycological study on some processed meat products exposed for sale in markets. *Assiut Vet. Med. J.*, 52(110): 121-131.
10. FAO (1986): Fisheries Technical Papers-T142. The production of fish meal and oil. Fisheries Industries Division, Food and Agriculture Organization of the United Nations, Rome, Italy. <http://www.fao.org/DOCREP/003/X6899E/X6899E00.HTM>
11. Frisvad, J.C. and Samson R.A. (2004): Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A guide to identification of food and air-borne terverticillate *Penicillia* and their mycotoxins. *Studies in Mycology* 49: 1-173.
12. Ghaly, A.E.; Dave, D.; Budge, S. and M.S. Brooks, (2010): Fish spoilage mechanisms and preservation techniques: Review. *Am. J. Applied Sci.*, 7: 846-864.
13. Henry, T.; Iwen, P.C. and Hinrichs, S.H. (2000): Identification of *Aspergillus* species using internal transcribed spacer regions 1 and 2. *J Clin. Microbiol*; 38: 1510-1515.
14. Huis in't Veld, J.H.J. (1996): Microbial and biochemical spoilage of foods: An overview. *Int. J. Food Microbiology*, 33: 1-18.
15. Hussein, M. A. M. (2008): Mycological Aspect of Fresh and Processed Meat Products with Special Respect to Proteolytic and Lipolytic Mold. Ph.D. Thesis, (Meat Hygiene) Fac. Vet. Med., Zagazig Univ., Egypt.
16. ISO (217-1-2:2008) "International Standards Organization EAST AFRICAN STANDARD: Microbiology of food and animal feeding stuffs — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination- Part 1-3: Specific rules for the preparation of meat and meat products, 2008.
17. Iwen, P.C.; Hinrichs, S.H. and Rupp, M.E. (2002): Utilization of the internal transcribed spacer regions as molecular targets to detect and identify human fungal pathogens. *Med Mycol*; 40: 87-109.
18. Jay, J.M.; Leossner, M.J. and Golden, D.V. (2005): *Modern food microbiology*. 7th Ed. Chapter 2, Taxonomy, Role, and Significance of Microorganisms in Foods. Pp, 13- 37, Springer Science.
19. Jooyandeh, H.; Amarjeet, K. and Minhas, K.S. (2009): Lipases in dairy industry: A review. *J Food Sci Technol*, 46(3): 181-189.
20. Korashy- Eman and Wahba-Nahed, M. (2008): lipolytic and Proteolytic activities of some fungi isolated from raw camel's milk. *Assiut Vet. Med. J.* 54: 118.
21. Makimura, K. (2001): Species identification system for dermatophytes based on the DNA sequences of nuclear ribosomal internal transcribed spacer 1. *Nippon Ishinkin Gakkai Zasshi*; 42:61-67.
22. Makimura, K.; Murayama, S.Y. and Yamaguchi H. (1994): Specific detection of *Aspergillus* and *Penicillium* species from respiratory specimens by polymerase chain reaction (PCR). *Jpn J Med Sci. Biol.*; 47:141-156.
23. Morshdy, A.E. M.A.; Hussien, M. A. M.; El-Abbasy M. T. and Elzwahery, R. R.M. (2015): Aflatoxins Residues in Some Meat Products.

- 2ndConference of Food Safety, Suez Canal University, Faculty of Veterinary Medicine, I: 90-95.
25. Nasser, L.A. (2002): "Mycological status of imported canned fish consumed in Saudia Arabia with special reference to proteolytic activity". Assuit Vet. Med. J., 47:125-135.
 26. Ouf, J. M.; Nagwa, I.M.K. and Shabana, E.S.E. (2010): Incidence of proteolytic and lipolytic moulds and yeasts in some ready to eat meat products. Assuit Vet. Med. J., 56(126):132-143.
 27. Pitt, J.I. and Hocking, A.D. (2009): Fungi and Food spoilage. 3rd Ed. Published by Springer Dordrecht Heidelberg London New York.
 28. Rohlf, F.J. (1998): NTSYSpc Numerical Taxonomy and Multivariate Analysis System Version 2.0 User Guide. Applied Biostatistics Inc., Setauket, New York. 37-34.
 29. Shaltout, F. A. (2002): Microbiological aspect of Semi-cooked chicken meat products. Benha Vet. Med., J., 13(2):15-26.
 30. Shaltout, F. A.; El-diasty, E.M. and Mohamed, M. S. (2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. 1st Scientific conference of food safety and Technology, pp. 79-89.
 31. Shaltout, F.A. and Salem, R.M. (2000): Molds, Aflatoxin B₁ and Ochratoxin A in frozen livers and meat products. Vet. Med. J. Giza. 48 (3): 341-346.
 32. Soliman, Z.I. and Shalaby, A. M. (2008): Occurrence of lipolytic and proteolytic fungi in locally smoked fish. Assiut Vet. Med. J., 54(117): 78-88.
 33. Subash, G.; Periasamy, A. and Azariah, H. (2005): Extracellular enzymatic activity profiles in fungi isolated from oil-reach environments. J. Mycoscience 46, (2):119-126.
 34. Sunitha, V.H.; Nirmala Devi, D. and Srinivas, C. (2013): Extracellular enzymatic activity of endophytic fungal strains isolated from medicinal plants. World Journal of Agricultural Sciences 9 (1): 1-9.
 35. Toldra, F. (2006): The role of muscle enzymes in dry cured meat products with different drying conditions. Trends in Food Sci. Techn., 17: 164-168.
 36. Yamakami, Y.; Hashimoto, A.; Tokimatsu, I. and Nasu, M. (1996): PCR detection of DNA specific for Aspergillus species in serum of patients with invasive aspergillosis. J Clin Microbiol.; 34:2464-2468.
 37. Zhao, J.; Kong, F.; Li, R.; Wang, X.; Wa, Z. and Wang, D. (2001): Identification of Aspergillus fumigatus and related species by nested PCR targeting ribosomal DNA internal transcribed spacer regions. J Clin Microbiol.; 9: 2261-/2266.

4/8/2019