

Comparison of *Helicobacter pylori* infection between pregnant women with hyperemesis gravidarum and controls

Fariba Nanbakhsh¹, Hamideh Mohaddesi^{*},² Fatemeh Bahadory¹, Javad Amirfakhrian³, Pouya Mazloomi³

1. Department of Obstetrics and Gynecology, Urmia University of Medical Sciences, Urmia, Iran
2. School of Nursing and Midwifery, Urmia University of Medical Sciences, Urmia, Iran
3. General Practitioner, Urmia University of Medical Sciences, Urmia, Iran
hmohaddesi.han@gmail.com

Abstract: A small percentage of pregnant women experience a severe form of nausea and vomiting called Hyperemesis gravidarum. To compare the rate of *H. pylori* infection between an Iranian samples of pregnant women who were suffering from hyperemesis gravidarum (HG) in their first trimester and control pregnant women without HG. In this cross-sectional study 30 pregnant women who suffered from HG and 30 controls were enrolled. *H. pylori* specific serum IgG was measured using ELISA method. IgG titers of more than 10, less than 10, and between 5 and 10 U/mL were considered as seropositive, seronegative, and suspicious for *H. pylori* infection, respectively. No significant difference was seen between the two groups regarding age, gestational age, and gravidity. Twenty-four patients (92.3%) in HG group had *H. pylori* infection, while this rate was only 7.7% (two patients) in control group ($P < 0.001$). No correlation was detected between IgG titers and either maternal age ($t=2.4$, $P = 0.12$) or parity ($\chi^2=0.2$, $P = 0.6$). There was significantly higher rate of *H. pylori* infection in HG pregnant women in their first trimester in comparison to controls..

[Fariba Nanbakhsh, Hamideh Mohaddesi, Fatemeh Bahadory, Javad Amirfakhrian, Pouya Mazloomi. **Comparison of *Helicobacter pylori* infection between pregnant women with hyperemesis gravidarum and controls.** *Life Sci J* 2012;9(4):1315-1318] (ISSN:1097-8135). <http://www.lifesciencesite.com>. 199

Keywords: Hyperemesis gravidarum; *helicobacter pylori*; pregnant women

1. Introduction

Nausea and vomiting is a major complaint in 70-80% of pregnancies. The typical onset is between 4 and 8 weeks and continues until 14-16 week of pregnancy (Cunningham et al, 2010). The condition is usually self limiting and peaks at around 9th gestational week. However, in up to 20% of cases, nausea and vomiting may continue until delivery. Nausea and vomiting occurs significantly more often in primigravidas and in women who were less educated, younger, non-smokers, obese, or those with a history of nausea and vomiting in a previous pregnancy (Jueckstock et al,2010).

A small percentage of pregnant women experience a severe form of nausea and vomiting called Hyperemesis gravidarum (HG). More than 59,000 pregnant women with HG are hospitalized in the US annually.

Asian populations tend to have higher incidence rates. For example, a Malaysian study identified 192 recorded cases (3.9%) out of 4,937 maternities (Fejzo et al, 2008)

HG is often associated with weight loss, ketonemia, ketonuria, dehydration, and acidosis from starvation, alkalosis from loss of hydrochloric acid, electrolyte disturbances and even hepatic and renal damage (Eliakim et al, 2000).

Estimate of incidence of HG varies widely depending on diagnostic criteria sand different study

populations. Some sources state a proportion of 0.5-2% of all pregnancies, whereas others provide an incidence of 0.3-1.5% of all live births (Aytac et al,2007.ACOG 2004).

HG can be associated with serious maternal and fetal morbidity such as Wernicke's encephalopathy, fetal growth restriction, and even maternal and fetal death (Bailit, 2005).

Although the pathophysiology for HG is not clear, many hypotheses have been proposed to explain the etiology and pathophysiology of HG including psychological factors, gastrointestinal tract dysfunction, endocrinologic changes, infections, immunological and metabolic causes, and anatomical factors. However, there is no single theory to provide an adequate explanation for all the manifestations of HG (Aytac et al, 2007. Karaca et al, 2004. Verberg, 2005. Niebyl, 2010).

A number of risk factors associated with HG have been reported including nulliparity, (Atanackovic, 2001) low maternal age(Eliakim et al, 2000. Kallen, 2003), multiple gestation (Eliakim et al, 2000. Basso and Olsen, 2001. Schiff et al, 2004), metabolic disturbances, trophoblastic disorders, fetal anomalies (Eliakim et al, 2000. Schiff et al, 2004), previous pregnancy complicated by HG,(Eliakim et al, 2000) female sex of the fetus (Basso and Olsen, 2001. Schiff et al, 2004), psychiatric conditions(Atanackovic, 2001., Swallow et al, 2004),

both high and low maternal pregnancy weight and lower socio-economic status (Eliakim et al, 2000. Schiff et al, 2004.. Rochelson et al, 2003). Smoking, on the other hand, has been associated with a reduced risk of HG (Kallen et al 2003. Schiff et al,2004).

Chronic infection with *Helicobacter pylori* (*H. pylori*) may also cause HG. *H. pylori* are a gram negative, spiral shaped, microaerophilic bacterium that colonizes the gastric mucosa of the human being and is associated with type B gastritis, peptic ulcer and cancers (Graham 2000). The prevalence rate is higher in developing countries especially among those with low socioeconomic status (Soll 1996). It is estimated that 50-65% of the world population is colonized with this bacterium (Jaakkimainen et al, 1999), but despite this high prevalence, only a proportion of these people are clinically manifested, the reason may be related to host defense factors(Mobley 1997).

Recently several studies have implicated *H. pylori* infection as a possible cause for HG. However, there are also reports that did not show any relationship between *H. pylori* infection and occurrence of HG. Nevertheless, no conclusive evidence has been adduced to suggest a direct causal relationship between *H. pylori* infection and HG. So, the role of *H. pylori* infection in HG is controversial (Karaca et al, 2004, Karadeniz et al 2006. Salimi-Khayati et al 2003. Erdem_ et al 2002. Wu et al 2000).

The purpose of the present study was to investigate any possible association between infection with *H. pylori* and HG. In the case of this association, eradication of *H. pylori* may help to treat HG in a group of Iranian pregnant women which is a problem in the first trimester of pregnancy.

2. Material and Methods

Sixty consecutive pregnant women in the first trimester of their pregnancy were entered into this cross-sectional study. Thirty women had HG (HG group) and 30 did not have HG (control group). The criteria for the presence of HG were pernicious vomiting (more than 3 times per day) without any obvious cause except of pregnancy, weight loss of more than 3 kg or 5% of body weight, and the presence of at least +1 ketonuria. Inclusion criteria for experimental group included having HG, age of 15-45 years, gestational age < 20 weeks and exclusion of other causes of vomiting such as hyperthyroidism, molar pregnancy, infectious diseases, psychological disorders, multiple pregnancies and gastrointestinal disorders. Thirty pregnant women in their first trimester were entered as control group. Inclusion criteria for control group were the same as for HG group except for symptoms of HG. The control group was selected among

asymptomatic pregnant women that were visited at the antenatal clinic of our hospital.

Venous blood was obtained at the first visit. *H. pylori* specific IgG antibody titer was measured using by enzyme-linked immune sorbent assay (ELISA) method. IgG titer less than 10 U/mL was considered negative and a titer more than 10 U/mL was interpreted as positive. IgG titer between 5 and 10 U/mL was considered as suspicious. Gestational age was determined using the date of the last menstrual period and was confirmed by ultrasonography. IgG measurement was done two times for all sera.

Statistical analysis was performed using SPSS software for Windows (ver. 10.0). Descriptive indices including mean, standard deviation (SD), frequency, and percentage were used to express data. For comparison of categorical variables, the chi-squared test and for continuous variables, the student t test was applied. Significance level was set at $P < 0.05$.

Informed consent was obtained from all participants. The study protocol was in accordance with the Helsinki Declaration and approved in ethics committee of Urmia University of Medical Sciences.

3. Results

Table 1 presents demographic characteristics of both groups. The age of the women in both groups ranged was 15-45 years. Mean (SD) age of HG group was 26.1 (± 2.5) years and in control group was 25.03 (± 2.8) years ($P = 0.4$). There was no significant statistical difference between the two groups regarding demographic variables.

Table 1. Demographic characteristics of patients and controls

Parameters	HG group	Controls	P value
Age, year	26.1 (± 2.5)	25.03 (± 2.8)	0.478
Gravidity	1.73	1.6	0.6
Parity	0.7	0.6	0.44
Gestational week	9.9	9.6	0.55
<i>H. Pylori</i> seropositivity	24 (92.3%)	2 (7.7%)	0.001

Table 2 presents *h. pylori* seropositivity in the studied groups. Twenty-four patients (92.3%) in HG group had *H. pylori* infection, while this rate was only 7.7% (two patients) in control group ($P < 0.001$; $\chi^2 = 32.8$).

Table 2. *H. pylori* seropositivity in HG and control groups

<i>H. pylori</i> serostatus	HG group	Controls	Total	P value
Positive	24 (92.3%)	2 (7.7%)	26	<
Negative	1 (5.9%)	16 (94.1%)	17	0.001
Suspicious	5 (29.4%)	12 (70.6%)	17	

In Table 3, mean (SD) values of H. pylori specific IgG antibody titer has been depicted. Mean (SD) IgG titer was 23.34 (± 13.9) U/mL (range, 4.7-62.6) in HG groups and was 6.09 (± 4.5) U/mL in control groups (range, 1.4-21.7).

One patient in HG group and 16 patients in control group had IgG titers of less than 5 U/mL. Borderline IgG titer (5-10 U/mL) was observed in 17 cases (5 patients (29.4%) in HG group vs. 12 cases (70.6%) in control group; $P < 0.001$).

We did not find any correlation between IgG titers and either maternal age ($t=2.4$, $P = 0.12$) or parity ($\chi^2=0.2$, $P = 0.6$).

There was no statistically significant difference in gestational age and gravidity between patients and controls ($\chi^2=0.44$, $p=0.55$).

Table 3. Comparison of mean (standard deviation) H. pylori specific IgG titer between HG patients and controls.

H. pylori specific IgG titer	Max.	Min.	Mean (\pm SD)	P value
HG group	62.6	4.7	23.4 (± 13.9)	< 0.001
Controls	21.7	1.4	6.09 (± 4.5)	
Total	62.60	1.4	14.71	

4. Discussions

It is estimated that H. pylori infection might be present in two-thirds of the world population. However, the pathogenic relationship between HG and H. pylori is not self evident because most of those infected with H. pylori do not complain of symptoms. In other words, the presence of H. pylori can be asymptomatic. Furthermore, the problems in diagnosis of H. pylori infection are more complicated during pregnancy since HG can mask an active H. pylori infection or HG may be worsened by superimposed H. pylori infection (Karadeniz et al 2006).

A possible explanation for an association between H. pylori infection and HG could be an increased accumulation of fluid and a displacement of intracellular and extracellular volumes which occur in the early phase of pregnancy as a result of increase in steroid hormones. These physiologic changes results in a change of gastric pH levels which could lead to the manifestation of a latent H. pylori infection in the gastrointestinal tract (Frigo et al 1998, Kocak et al 1999). The results of this study show that the seropositivity of H. pylori in the HG group was significantly higher than in control group. Recently, it was shown that H. pylori seropositivity was significantly high in pregnant women who suffer from HG. Our findings are in accordance with the previous studies (Karaca et al 2004, Frigo et al 1998,

Kocak et al 1999). However, some recent studies could not find such association. Thus this is one of the controversial issues in obstetric care (Jacobson et al 2003, Berker et al 2003). For example, Berker et al. reported insignificant difference between HG patients (70%) and controls (61.3%) regarding the presence of H. pylori infection using H. pylori specific IgG. Similar results were noted by Lee et al. (Mobley 1997) who demonstrated the prevalence of H. pylori infection in HG patients to be 65%, while this figure was 66.7% in controls.

Frigo et al. reported statistically significant differences between HG patients and asymptomatic ones regarding H. pylori infection (90.5% vs. 46.5%) (Frigo et al 1998). Kocak et al also found similar results (92% in HG patients vs. 45% in controls) (Kocak et al 1999). Salimi-Khayati et al. revealed significantly higher H.pylori seropositivity in HG patients (88.9%) than in control ones (40.7%) Salimi-Khayati et al 2003).

In a total of 12 studies reviewed regarding H. pylori infection in HG patients, 11 relied on the serologic assay of H. pylori infection, and the great majority of them showed significantly increased infection rate in HG patients. The authors of three studies were not able to demonstrate any meaningful difference in H. pylori infection rates between HG patients and controls (Jacobson et al 2003, Berker et al 2003).

Several results were not able to demonstrate any significant differences between HG patients and asymptomatic ones as regards H. pylori IgG antibody seropositivity (Jacobson et al 2003).

Aytac et al. did not find any significant difference between HG patients and control ones (41.1% vs. 40%) suggesting no causal relationship between H. pylori infection and HG (Aytac et al 2007). Karadeniz et al. reported the prevalence of H. pylori IgG antibody was 67.7% (21 of 31) in the patients with HG and 79.3% (23 of 29) in controls ($P = 0.31$). There was no statistically significant difference in study and control groups for H. pylori IgG Ab and H. pylori SA in that study. (Karadeniz et al 2006).

Two recent studies found no association between HG and Hp seropositivity, one conducted in two US populations with disparate Hp seropositivity and the other by Berker et al from Turkey (Jacobson et al 2003, Berker et al 2003).

5. Conclusion:

This study suggests an association between H. pylori infection and hyperemesis gravidarum. Serologic test, moreover, being a non-invasive, cheap and easy to use, we recommend its routine use in patients with hyperemesis gravidarum as well as in

women who desire to become pregnant in the near future.

Acknowledgments:

The authors wish to thank the colleagues and patients that were kindly engaged and collaborated with us throughout this study.

Corresponding Author:

Dr. Hamide Mohaddesi
Department of Obstetrics and Gynecology, Urmia,
University of Medical Sciences, Urmia, Iran
E-mail: hmohaddesi.han@gmail.com

References

- Cunningham G, Gant NF, Kenneth JL. William's Obstetrics. 21st ed. McGraw-Hill Company, Philadelphia, 2001.
- Jueckstock JK, Kaestner R, Mylonas I. Managing hyperemesis gravidarum: a multimodal challenge. *BMC Med* 2010; 8: 46.
- Fejzo MS, Ingles SA, Wilson M. High prevalence of severe nausea and vomiting of pregnancy and hyperemesis gravidarum among relatives of affected individuals. *Eur J Obstet Gynecol Reprod Biol* 2008; 141: 13- 17.
- Eliakim R, Abulafia O, Sherer DM. Hyperemesis gravidarum: a current review. *Am J Prenatal* 2000;17: 207-18.
- Aytac S, Türkay C, Kanbay M. Helicobacter pylori stool antigen assay in hyperemesis gravidarum: a risk factor for hyperemesis gravidarum or not? *Dig Dis Sci* 2007; 52: 2840-83.
- ACOG (American College of Obstetrics and Gynecology): Practice bulletin: nausea and vomiting of pregnancy. *Obstet Gynecol* 2004; 103: 803-14.
- Bailit JL. Hyperemesis gravidarum: epidemiologic findings from a large cohort. *Am J Obstet Gynecol* 2005;193:811-14.
- Karaca C, Güler N, Yazar A. Is lower socio-economic status a risk factor for Helicobacter pylori infection in pregnant women with hyperemesis gravidarum? *Turk J Gastroenterol* 2004; 15: 86-9.
- Verberg MF, Gillott DJ, Al-Fardan N. Hyperemesis gravidarum, a literature review. *Hum Reprod Update*. 2005; 11: 527-39.
- Niebyl MD. Clinical practice. Nausea and vomiting in Pregnancy. *N Engl J Med* 2010;363: 1544-550
- Atanackovic G, Wolpin J, Koren G. Determinants of the need for hospital care among women with nausea and vomiting of pregnancy. *Clin Invest Med* 2001; 24:90- 93.
- Kallen B, Lundberg G, Aberg A. Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. *Acta Obstet Gynecol Scand* 2003, 82,916-20.
- Basso O, Olsen J. Sex ratio and twinning in women with hyperemesis or pre-eclampsia. *Epidemiology* 2001;12: 747-9.
- Schiff MA, Reed SD, Daling JR. The sex ratio of pregnancies complicated by hospitalisation for hyperemesis gravidarum. *BJOG* 2004; 111: 27- 30.
- Swallow BL, Lindow SW, Masson EA. Psychological health in early pregnancy: relationship with nausea and vomiting. *J Obstet Gynaecol* 2004; 24: 28 -32.
- Rochelson B, Vohra N, Darvishzadeh J. Low prepregnancy ideal weight: height ratio in women with hyperemesis gravidarum. *J Reprod Med* 2003; 48: 422- 24.
- Graham DY. Helicobacter pylori infection is the primary cause of gastric cancer. *J Gastroenterol* 2000; 35: 90-97.
- Soll AH. Helicobacter pylori induced gastritis. In: Bennet JC, Plum F, eds. Cecil Textbook of Medicine, (20th edn). Philadelphia: WB Saunders; 1996.
- Jaakkimainen RL, Boyle E, Tudiver F. Is Helicobacter pylori associated with non-ulcer dyspepsia and will eradication improve symptoms? A Meta-analysis. *BMJ* 1999;319: 1040 -44.
- Mobley H. Helicobacter pylori factors associated with disease development. *Gastroenterology* 1997; 113: 521- 8.
- Karadeniz RS, Ozdegirmenci O, Altay MM. Helicobacter pylori seropositivity and stool antigen in patients with hyperemesis gravidarum. *Infect Dis Obstet Gynecol* 2006;2006:73073.
- Salimi-Khayati A, Sharami H, Mansour-Ghanaei F. Helicobacter pylori aeropositivity and the incidence of hyperemesis gravidarum. *Med Sci Monit*. 2003;9: CR12-5.
- Erdem A, Arslan M, Erdem M. Detection of Helicobacter pylori seropositivity in hyperemesis gravidarum and correlation with symptoms. *Am J Perinatol* 2002; 19: 87- 92.
- Wu CY, Tseng JJ, Chou MM. Correlation between Helicobacter pylori infection and gastrointestinal symptoms in pregnancy. *Adv Therapy* 2000; 17: 152- 8.
- Frigo P, Lang C, Reisenberger K. Hyperemesis gravidarum associated with Helicobacter pylori seropositivity. *Obstet Gynecol* 1998; 91: 615-17.
- Kocak I, Akcan Y, Ustun C. Helicobacter pylori seropositivity in patients with hyperemesis gravidarum. *Int J Gynecol Obstet* 1999; 66: 251-4.
- Jacobson GF, Autry AM, Somer-Shely TL. Helicobacter pylori seropositivity and hyperemesis gravidarum. *J Reprod Med* 2003; 48: 578 -82.
- Berker B, Soylemez F, Cengiz SD. Serologic assay of Helicobacter pylori infection. Is it useful in hyperemesis gravidarum? *J Reprod Med* 2003; 48: 80 -2.
- Lanciers S, Despinasse B, Mehta DI. Increased susceptibility to Helicobacter pylori infection in pregnancy. *Infect Dis Obstet Gynecol* 1999;7: 195-8.

9/29/2012