

Biological Activities of Characterized Isolates of n-Hexane Extract of Azadirachta Indica A.Juss (Neem) Leaves

A.Akpuaka¹, M.M. Ekwenchi¹, D.A. Dashak¹, A. Dildar²

1. Department of Chemistry, Faculty of Natural Sciences, University of Jos, Jos, Plateau State, Nigeria.

2. Department of Chemistry, Foreman Christian College (A Chartered University), Lahore, Pakistan.

aakpuaka33@gmail.com

Abstract: The antimicrobial activities of n-Hexane extract and 5 Column Chromatography fractions, of Azadirachta indica A. juss (Neem) leaves, showed antimicrobial activities against human pathogenic bacteria (*Salmonella typhi*) and yeast fungus (*Candida albicans*). Antimicrobial properties of Azadirachta indica fractions were tested using ditchwell diffusion method. Analysis of the data projected that upon bioassay with n-Hexane neem leaves extract and 5 fractions of Column Chromatography zone of inhibition for *Salmonella typhi* observed was 17mm, 12mm, 5mm, 3mm and 1mm respectively. While for the yeast fungus *Candida albicans* the zones of inhibition, seen were 28mm, 25mm, 20mm, 3mm, 21mm and 20mm for Azadirachta indica, respectively. Results were compared to conventional drugs. GC/MS identified 45 bioactive compounds in the n-Hexane extract of Azadirachta indica leaves out of which 33 have antifungal activity. Conclusively based on the data analysis it can be said Azadirachta indica A. juss leaves extract have biological activity as good as the conventional drugs against such microorganisms.

[Akpuaka A., Ekwenchi M M., Dashak D A., Dildar A. **Biological Activities of Characterized Isolates of n-Hexane Extract of Azadirachta Indica A.Juss (Neem) Leaves.** *N Y Sci J* 2013;6(6):119-124]. (ISSN: 1554-0200). <http://www.sciencepub.net/newyork>. 17

Keywords: Azadirachta indica A. Juss; n-Hexane neem Leaf extract; Antifungal activity; Antibacterial activity; Zone of Inhibition;

1. Introduction

Salmonella is a genus of rod-shaped, Gram negative non-spore forming, non predominantly motile enterobacteria with 2 to 5µm and flagella that grade in all direction (Clark M.A. et al., 1987) (Ryan K.J. et al., 2004). They cause typhoid fever, paratyphoid fever, and food borne illness.

Candida albicans is a diploid fungus that grows both as yeast and filamentous cells and a causal agent of opportunistic oral and genital infection in humans (Ryan K.J. et al., 2004).

It is a parasitic fungus that can infect mouth or the skin or the intestines or the vagina and causes candidiasis. Systemic fungal infection (fungemias) causes morbidity in immunocompromised patients (e.g AIDS, Cancer chemotherapy etc). *Candida albicans* is a commensal and a constituent of the normal gut flora of 80% of the human.

Azadirachta indica belongs to the Meliaceae family and is a drought resistant tropical to

Leaves of *Azadirachta indica* A. Juss were collected from the National Research Institute of Chemical Technology, Zaria – Kaduna State Nigeria. Authenticated sample is deposited in the Department's herbarium, of Plant Science Technology, University of Jos, Jos, Plateau State, Nigeria. The washed leaves of Neem were dried in an electric oven at 30°C for 4 days and powdered with an electric blender.

500g of *Azadirachta indica* A. Juss Leaves were powdered and extracted for 4 days with n-Hexane

subtropical tree. For centuries, it has served mankind as “Panacea for all diseases”. It is antifungal (Ramasany H, et al., 2010), antiviral (Badam L, et al 1999), contraceptive and sedative (Medicinal Properties of Neem, see references), antioxidant (Syeda et al., 2011), antiretroviral, antimicrobial (Sewanu, et al., 2012), anti-inflammatory, active against resistant Gonorrhoea (Okeie, et al., 2009), antibacterial (Yogeswari, et al., 2012), anticancer (Raldary, et al., 2012), antitumor (Hsonuna, et al., 2011), antimalaria (Pavlović, et al., 2011), antidiabetic (Gholamreza, et al., 2012).

Analysis confirmed presence of phytochemicals like Alkaloids, Steroids, Tannins, Terpenoids, Flavonoids, Polyphenols and Saponins are found to be present in Neem Leaves. Current study is carried out to establish the biological activity of *Azadirachta indica* leaves.

2. Materials and Methods

with constant shaking. Extract was dried to constant weight in a vacuum hood. Extract was applied to Column for Column Chromatography using 5 solvent systems. **1.** 100% n-Hexane **2.** 95% n-Hexane + 5% Benzene **3.** 85% Benzene + 15% n-Hexane **4.** 20% Benzene + 20% Diethylether + 60% Methanol **5.** 100% Methanol respectively.

The organisms for present study were clinical isolates of *Salmonella typhi* from the National Veterinary Research Institute (NVIR), Vom, Plateau state, Nigeria. *Candida albicans* were clinical isolates

from the Department of Pathology, ALLAMA IQBAL Medical University Lahore, Pakistan.

n-Hexane extract and its 5 Column Chromatography fractions of *Azadirachta indica* were bioassayed against the two organisms separately. The Potatoe Dextrose Agar plates were each swabbed on the surface with the respective organism. Using cork borer, wells were carved and 1 mg/ml of the n-Hexane neem leaves extract and 1mg/ml of each of the 5 Column Chromatography fractions were applied to the wells respectively. Antibacterial positive control was Chlorophenicol 1mg/ml and distill water was the negative control. Antifungal positive control was Fluconazol 1mg/ml and for negative control distill water was employed. The plates in triplicates were incubated at 28°C for 48hours. The diameter of zones of inhibition in millimetre of the test organisms were noted.

The GC-MS analysis of n-Hexane *Azadirachta indica* leaves extract and 5 Column Chromatography fractions were carried out using Agilent Technologies 7890A GC system. The detector was Agilent Technologies 5975C inert MSD with Triple-Axis Detector, with column (Polysiloxanes) 30m × 0.25m

fused capillary silica tubing. Software adopted to handle mass spectra and chromatograms was National Institute of Standards and Technology MS. 2005 Library. The following temperature protocol was followed for GC/MS detection. Injection port temperature was 200°C and Helium flow rate was 1ml/min. Oven temperature was programmed from 50°C with an increase of 8°C/min to 300°C and this temperature was held for 9 minutes. The ionization voltage was 70 eV. The samples were injected in splitless mode and mass spectral scan range was set at 45-500(MHZ). The GC/MS characterized the bioactive components of the n-Hexane extract Column Chromatography fractions. The fragmentation pattern of the mass spectra were compared (head to tail) with those stored in the NIST Library. The total running time for GC was 36 minutes.

3. Results

The Results are presented as follows:

Table 1: indicates the zones of inhibition of the bacteria, while Table 2: shows the zones of inhibition for the fungus.

Table 1. Zones of Inhibition of *Salmonella typhi* against *Azadirachta indica* A. juss n-Hexane leaves extract and Column Chromatography fractions

S/N	Solvent System	Column Chromatography Fractions	Zone of inhibition (mm)
0.	n-Hexane Neem extract	Extract	17
1.	n-Hexane fraction	(CC F1)	12
2.	95% n-Hexane + 5% Benzene	(CC F2)	5
3.	85% n-Hexane + 15% Benzene	(CC F3)	3
4.	20% Benzene + 20% Diethylether + 60% Methanol	(CC F4)	11
5.	100% Methanol	(CC F5)	No Inhibition

Table 2. Zones of Inhibition of *Candida albicans* against *Azadirachta indica* A. juss leaves with n-Hexane extract and Column Chromatography fractions

S/N	Solvent System	Column Chromatography Fractions	Zone of inhibition (mm)
0.	n-Hexane Neem extract		28
1.	n-Hexane fraction	(CC F1)	25
2.	95% n-Hexane + 5% Benzene	(CC F2)	20
3.	85% n-Hexane + 15% Benzene	(CC F3)	3
4.	20% Benzene + 20% Diethylether + 60% Methanol	(CC F4)	21
5.	100% Methanol	(CC F5)	20

Table 3. shows the Gas Chromatography and Mass Spectroscopy characterized components.

Table 3: GC-MS Identified components of the *Azadirachta indica* A. juss Leaves extract (Compounds are listed in ascending order of Retention Time)

S/N	RT	Molecular Formula	MW	Name of Compound	Activity
1. ****	4.237	C8H10	106	p-Xylene	Antifungal, Antioxidant and antimicrobial (7)
2. ****	4.250	C9H20	128	Nonane	Antifungal and Antibacterial (12)
3. ****	4.262	C8H10	106	o-Xylene	Antifungal, Antioxidant and antimicrobial (7)
4. ****	4.806	C8H9NO2	151	Oxime-, methoxy-phenyl-	Antifungal, Antibacterial, Anticancer and Antitumor (10)
5. ****	5.845	C10H22	142	Decane	Antifungal and Antibacterial (13)
6. ****	6.389	C10H14	134	Benzene,1-methyl-2-(1-methylethyl)-	Potent antifungal Antioxidant activity (14)
7. ****	6.495	C7H8O	108	Benzyl Alcohol	Potent antifungal Active against yeasts, mold and more active against Gram-positive, than Gram-Negative Bacteria. (15)
8. ****	7.709	C9H18O	142	Nonanal	Antifungal Moderate Antibacterial activity against Gram +ve and Gram- bacteria (16)
9. ****	9.179	C7H6O2	122	Benzoic Acid	Antifungal (17)
10. ****	9.391	C12H26	170	Dodecane	Enhances antifungal activity (18)
11. ****	12.213	C10H12O2	164	Phenol, 2-methoxy, 3-(3-propenyl)-	Antifungal, High level of antimicrobial activity against fish pathogens, antioxidant, and disinfectant. (3)
12. ****	12.769	C14H30	198	Tetradecane	Antifungal and Antibacterial (19)
13. ****	13.376	C10H18O3	186	Nonanoic acid, 9-oxo-, methyl ester	Potent antifungal, Antioxidant, Potent Antimicrobial (6)
14. ****	15.403	C15H24	204	.gamma.-Elemene	Antifungal, Antioxidant, and Biocidal activity (20)
15. ****	15.703	C15H24O	220	1H-Cycloprop(e)azulen-7-ol, decahydro-1,1,7-trimethyl-4-methylene-, {1aR-(1aalpha,4aalpha,7beta,7abeta,7balpha)}-	Antifungal, Insecticidal and larvicidal, agent (21)
16. ****	15.829	C16H34	226	Hexadecane	Antifungal, Antibacterial, antioxidant activity, (9;11)
17. ****	18.568	C19H40	268	Nonadecane	Cytotoxic effect, antimicrobial (11)
18. ****	18.918	C17H34O2	270	2(4H)-Benzofuranone, 5,6,7,7 ^a -tetrahydro-4,4,7a-trimethyl-	Antifungal, Antialgal effect, antioxidant, antibacterial activity (22;23)
19. ****	19.181	C18H36O	268	2-Pentadecanone, 6,10,14-trimethyl-	Antibacterial activity against Gram +ve and Gram-ve bacteria (16)
20. ****	19.537	C16H22O4	278	Diisobutyl phthalate	Antifungal, Antibacterial, Antiviral and Antioxidant activities (24)
21. ****	20.188	C17H34O2	270	Pentadecanoic acid, 14-methyl-, methyl ester	Antifungal, Antimicrobial (25)
22. ****	20.194	C17H34O2	270	Hexadecanoic acid, methyl ester	Antifungal, Antioxidant, hypocholesterolemic nematocide, pesticide, anti-androgenic flavour, haemolytic, 5-Alpha reductase inhibitor, potent antimicrobial activity. (26)
23. ****	20.463	C18H28O3	292	Benzenepropanoic acid, 3,5-bis(1,1-dimethylethyl-4-hydroxy-, methyl) ester	Antifungal, antioxidant, (25)
24. ****	20.463	C20H40O	296	1-Hexadecen-3-ol, 3,5,11,15-tetramethyl	Antifungal, Antiviral, Anti-inflammatory, Resistant gonorrhoea, (8)
25. ****	20.670	C16H32O2	256	n-Hexadecanoic acid	Antifungal, Antioxidant, hypocholesterolemic, nematocide, anti-androgenic flavour, haemolytic, 5-Alpha reductase inhibitor, potent antimicrobial agent, antimalarial and antifungal. (26; 27;11)
26. ****	20.713	C16H22O4	278	Dibutyl phthalate	Antifungal, Antimicrobial agent, antimalarial and antifungal (28)
27. ****	21.064	C12H25I	296	1-iodo-2-methylundecane	Antifungal, antioxidant and antimicrobial agents. (29) Insecticidal (30)
28. ****	21.064	C20H42	268	Eicosane	Antifungal, antibacterial, antitumor and cytotoxic effects (11)

29.	22.277	C19H36O2	296	9-Octadecenoic acid (Z)-methyl ester	Antioxidant activity, Anticarcinogenic,-exist in human blood and urine and serve as endogenous peroxisome proliferator-activated receptor ligand, dermatitogenic flavour (6;26)
30.	22.290	C19H32O2	292	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	Anticancer,Antimicrobial, Antioxidant and Hypercholesterolemic (31)
31. ****	22.434	C20H40O	296	Phytol	Antifungal Active against Salmonella typhi, Resistant Gonorrhoea, Joint dislocation, Headache, Hernia and is a Stimulant and anti malaria (26;6;8)
32. ****	22.565	C19H38O2	298	Octadecanoic acid, methyl ester	Potent Antifungal, antimicrobial, antibacterial and at low PH. (32)
33. ****	22.959	C18H36O2	284	Octadecanoic acid	Antifungal, antitumor activity, antibacterial (11;32)
34. ****	23.356	C22H48	310	Docosane	Antibacterial activity (33)
35. ****	26.975	C16H22O4	278	1,2-Benzenedicarboxylic acid, heptylmethyl ester	Active. New Compound
36.	26.975	C16H22O4	278	1,2-Benzenedicarboxylic acid, ethylhexyl ester	New Compound. Activity not tested yet.
37.	26.981	C16H22O4	278	1,2-Benzenedicarboxylic acid, mono(n-octyl) ester	New compound. Inactive.
38. ****	27.013	C16H22O4	278	1,2-Benzenedicarboxylic acid, mono(2-ethylhexyl) ester	Antifungal, anti retroviral, anti tumor,anti diabetic anti cancer,antioxioidant, anti scabies anti inflammatory, potent antimicrobial agent (6;34;35)
39.	28.301	C27H56	380	Heptacosane	Antioxidant activity (36)
40.	30.059	C29H60	408	Nonacosane	Antibacterial activity (37)
41.	30.872	C30H62	422	Triacotane	Cytotoxic effect, antimicrobial (11)
42. ****	31.704	C31H64	436	Hentriacontane	Antifugal against fungal spores germination, Antioxidant, antitumor activity and antibacterial (38)
43.	33.537	C29H48O	412	Cholesta-22,24-dien-5-ol,4,4-dimethyl-	Antibacterial, trypanocidal activity, (39)
44. ****	33.618	C33H68	464	Trtriacontane	Potent antifungal, Antibacterial, antioxidant activity (40)
45.	34.291	C29H50O	414	Gamma-Sitosterol	Antioxidant, antibacterial and prophylactic activities (41)

Key: **** = Antifungal components

4. Discussions

In the current study, the bioassay of n-Hexane neem leaves extract and 5 fractions against *Salmonella typhi* and *Candida albicans* shows activity. The antibacterial activity is moderate as inhibition zones are 17mm, 12mm, 5mm, 3mm and 1mm respectively, (Table 1:) The antifungal activity of the *Azadirachta indica* leaves is very significant, 28mm, 25mm, 20mm, 21mm and 20mm as seen in (Table 2:) GC-MS analysis identified total of 45 compounds in *Azadirachta indica* n-Hexane leaves extract viz: p-Xylene; Nonane; o-Xylene; Oxime-,methoxy-phenyl-; Decane; Benzene,1-methyl-2-(1-methylethyl)-; Benzyl Alcohol; Nonanal; Benzoic acid; Dodecane; Phenol, 2-methoxy, 3-(3-propenyl)-; Tetradecane; Nonanoic acid, 9-oxo- methyl ester; .gamma.-Elemene; 1H-Cycloprop(e)azulen-7-ol, decahydro-1,1,7-trimethyl-4-methylene-, {1aR-(1aalpha,4aalpaha,7beta,7abeta,7balpaha)}-; Hexadecane; Nonadecane; 2(4H)-Benzofuranone, 5,6,7,7^a-tetrahydro-4,4,7a-trimethyl-; 2-Pentadecanone 6,10,14-trimethyl-; Diisobutyl phthalate; Pentadecanoic acid, 14-methyl-,methyl

ester; Hexadecanoic acid, methyl ester; Benzenepropanoic acid, 3,5-bis(1,1-dimethylethyl-4-hydroxy-, methyl) ester; 1-Hexadecan-3-ol,3,5,11,15-tetramethyl; n-Hexadecanoic acid; Dibutyl phthalate; 1-iodo-2-methylundecane; Eicosane; 9-Octadecenoic acid (Z)-methyl ester; 9,12,15-Octadecatrienoic acid, methyl ester (Z,Z,Z)-; Phytol; Octadecanoic acid, methyl ester; Octadecanoic acid; Docosane; 1,2-Benzenedicarboxylic acid, methylheptyl ester; 1,2-Benzenedicarboxylic acid, methylheptyl ester; 1,2-Benzenedicarboxylic acid, mono(n-octyl) ester; 1,2-Benzenedicarboxylic acid, mono(2-ethylhexyl) ester; Heptacosane; Nonacosane; Triacotane; Hentriacontane; Cholesta-22,24-dien-5-ol,4,4-dimethyl-; Trtriacontane; and Gamma-Sitosterol. Their various activities are also mentioned in Table 3: Interestingly these components found in the n-Hexane extract differ so much from the components found in the Ethanolic extract of *Azadirachta indica* cited in (42), in which most of their components were Alkaloids while ours were mainly Terpenoids and Diterpenoids along with other non-polar components.

Acknowledgement:

I am grateful to the Department of Chemistry, University of Jos for their Laboratory facilities extended to me for carrying out my research work.

I thank the Chairman and all staff of the Department of Chemistry, Foreman Christian College Lahore, Pakistan for their Instrumentation facilities extended to me for this research work.

Corresponding Author:

Azra Akpuaka
Department of Chemistry
University of Jos, Jos, Nigeria
aakpuaka33@gmail.com

References

- Clark M.A., Barret E.L., (June, 1987)
- Ryan K.J. and Ray C.G. (2004). Sherris Medical Microbiology (4th ed.). Mc Graw Hill. 362-8. ISBN 0-8385-8529-9
- Ramasany, H., Man, C.K., Ju, S.K., Chellam, B., Sundaram, J., and Moon, S.H., (2010). Identification and Antimicrobial Activity of combined extract from *Azadirachta indica* and *Ocimum Sanctum*. The Israeli Journal of Aquaculture-Baidgeh, 62(2), 85-95.
- Badam L., Joshi S.P., (1999). "Invitro antifungal activity of neem (*Azadirachta indica* A. Juss) leaf extract against group B coxsackie viruses" J. Commun Dis., 31: 79-90.
- Medicinal Properties of Neem (http://www.infinityfoundation.com/mandala/t_es/t_es_agraw_neem.htm)
- Syeda, F.A, Habib-Ur- Rehman, Choudahry, M.I., and Atta-Ur-Rahman. (2011). Gas Chromatography-Mass Spectrometry (GC-MS) analysis of petroleum ether extract (oil) and bioassays of crude extract of *Iris germanica*. International Journal of Genetics and Molecular Biology, 3(7), 95-100.
- Sewanu, S.O., Opoku, A.R., Oyedeji, A.O (2012). The chemical Composition, antimicrobial and antioxidant properties of the essential oils of *Tulbaghia violacea* Hary and *Eucalyptus grandis* W.Hill ex Maiden. En US. University of Zululand South Africa.
- Okeie, W., Ogunlesi, M., Ofor, E., and Osibote, E.A.S (2009). Analysis of essential oil constituents in hydro-distillates of *calotropis procera* (oil). R.Br Res. Journal of phytochemistry 3, 44-53.
- Yogeswari, S., Ramalakshmi, S.N., and Muthu, J.M. (2012) identification and comparative studies of different volatile fractions from *Monochaetia kansensis* by GC-MS. Global Journal of Pharmacology, 6(2): 65-71.
- Rahdary, A.A. and Sobati, Z. (2012). Chemical composition of essential oil, antibacterial activity and brine shrimp lethality of ethanol extracts from *Sedum pallidum*. Journal of Medicinal Plants Research, 6(16), 3105-3109.
- Hsouna, A.B., Trigie, M., Mansour, R.B., Jarraya, R.M., Damak, M., and Jaoua, S., (2011). Chemical composition, cytotoxicity effect and antimicrobial activity of *Ceratonia silisqua* essential oil with preservative effects against *listeria* inoculated in minced beef meat. International Journal of Food Microbiology, 148(1), 66-72.
- Pavlović, M., Petrović, S., Milenković, M., Couladis, M., Tzakou, O., and Niketić, M. (2011). Chemical composition and antimicrobial activity of *Athriscus nemorosa* root essential oil. J. Nat. Prod. Commun., 6(2):271-273.
- Gholamreza, A., Mohamad, J. and Ehsan, S. (2012). Antimicrobial activity and chemical composition of essential oil from the seeds of *Artemisia aucheri* Boiss. J. Nat Pharm. Prod., 7(1):11-15.
- Harami, M., Adamu, E.O., Ekanem and Suleiman, B. (2010). Identification of Essential oil components from *Nigella sativa* seed by Gas Chromatography-mass Spectroscopy. Pak. J.Nutr., 9(10):966-967.
- David, P.E. and Patrick, J.C. (2012). Antimicrobial preservatives Part Two: Choosing a preservative. GlaxoSmithKline Pharmaceuticals Callum Consultancy.
- Nurettin, Y., Canan, G., Osman, U., Ahmet, Y., Serdar, U., Kamil, C., and Salih, T., (2006). Composition and antimicrobial activities of volatile components of *Minuaritia meyeri*. Turk. J. Chem., 30; 70-76.
- López, A., Ming, D.S., Towers, G.H., (2002). Antifungal activity of benzoic acid derivatives from *Piper lanceaefolium*. J. Nat. Prod., 65(1):62-64.
- Cheila, D.O.S., Fabricio, M.C., Fabricio, A.O., Evandro, P., Carmen, B.B.F., Susana, M.W.S., Maria, L.S. (2012). Antimicrobial activity [2methacryloyloxy)ethyl]trimethylammonium chloride against *Candida* spp,1-4.
- Ozdemir, G., Karabay, N.U., Dalay, M.C. Pazarbasi, B. (2004). Antibacterial activity of volatile component and various extracts of *Spirulina platensis*. Journal of Phytother Res, 18(9):744-757.
- Gurdip, S. and Palanisamy, M. (2006). Chemical constituents, antifungal and antioxidative effects of *ajwain* essential oil and its acetone extract. Journal of Agricultural and food chemistry, 54(1):174-81.

21. Madhumitha, G., Rajakumar, G., Roopan, S.M., Rahuman, A.A., Priya, K.M., Khan, F.R., Khanna, V.G., Velayutham, k., Jayaseelan, C., Kamarai, C., Elango, G., (2011). Acaricidal, Insecticidal, and larvicidal efficacy of fruit peel aqueous extract of *Annona Squamosa* and its compounds against blood-feeding parasites.
22. Akhari M., Batooli, H., and Kashi, F.J.,(2012). Composition of essential oil and biological activity of extracts of *Viola odoranta* L. from central Iran. *Journal of Nat. Prod. Res*, 26(9): 802-9.
23. Yang, W.D., Liu, Y.R., Liu, J.S., Liu, Z., (2008). Inhibitory effects and chemical basis of *Eucalyptus orelliana* wood meals on the growth of *Alexanrium tamarense*. *Journal of Huan Jing ke xue*, 29(8):2296-2301.
24. Yanping, H., Suhua, H., Jincheng, W.U., Shunquan, L. (2010). Identification of essential oils from the leaves of 11 species of *Eriobotrya*, 4379-4382.
25. Bashir, A. Ibrar, K. Shumaila, B. and Sadiq, Azam. (2012). Chemical composition and antifungal, phytotoxic, brine shrimp cytotoxicity, insecticidal, and antibacterial activities of the essential oils of *Acacia modesta*. *Journal of Medicinal plants Research*, 6(31), 4653-4659.
26. Hema, R., Kumaravel, S., and Alagusundaram. (2011). GC/MS Determination of Bioactive components of *Murraya koenigii*. *Journal of American Science*, 7(1).
27. Pietro, Z., Maurizio, S., Maurizio, B., Antonella, M., Sergio, R., Carmen, F. and Felice, S., (2010). Essential oil composition of stems and fruits of *Caralluma europaea* N.E.Br. (*Apocynaceae*). *Molecules*, 15, 627-638.
28. Elija, K., Vaishali, B., Adsul, M.K., Deshpande, N.R. and Kashalkar, R.V. (2012). Antibacterial activity of Dibutyl Phthalate : A secondary metabolite isolated from *Ipomoea carnea* stem. *Journal of Pharmacy Research*, 5, No 1.
29. Laila, S.A.H, Mohammad, A.H., Afaf, Mohammed, W. Qasim, A., Jamal N.A. (2012). Gas chromatography-mass spectrometry analysis of different organic crude extracts from the local medicinal plant of *thymus vulgaris* L.
30. Sahayaraj, K. and Ravi, C., (2008). Preliminary phytochemistry of *ipomea carnea* jacq and *vitex negundo* linn leaves. *Int. J. Chem. Sci*, 6-1.
31. Praveen P.K., Kumaravel, S., and Lalitha, C. (2010). Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. *African Journal of Biochemistry Research*, 4(7), 191-195.
32. Gehan, M.A., Hanan, A.E., Hassan, A.H.I., and Okbah, M.A., (2009). Marine natural products and their potential applications as anti-infective agents. *World Sciences Journal*, 7(7): 872-880.
33. Uma, B., and Parvathavarthini, R. (2010). Antibacterial effect of Hexane Extract of sea urchin, *Temnopleurus alexandri* (Bell, 1884). *International Journal of Pharm. Tech. Research CODEN (USA)*, 2(3) 1677-1680.
34. Balachandran, C., Lakshmi, R.S., Duraipandiyar, V., and Ignacimuthu S., (2012). Antimicrobial activity of *Streptomyces* sp. (ERI-CPDA-1) isolated from oil contaminated soil from Chennai, India. *Bioresource Technology*, 129.
35. Bagavathi, P.E., and Ramasamy, N., (2012). GC-MS analysis of phytochemicals in the ethanol extract of *Polygonum chinense* L. *Pharmacognosy Research*, 14, 11-14.
36. Fethi B.J.K., Mohamed, A.M., Samia, A.M., Karima, B.S., Ahmed, N.H., and Zine, M., (2012). Antioxidant activities of essential oil from *Cotula coronopifolia* L. growing in Tunisia.
37. Rani, P.S. and Agrawal, R. (2006). Volatile compounds of therapeutic importance produced by *leuconostoc paramesenteroides*, a native laboratory isolate.
38. Atolani, O., Olatunji, A.G., Adeyemi, O.S., and Fayemi O.S., (2009). Antioxidant and Antimicrobial Activity of Cuticular Wax from *kigelia africana*. *FABAD J. Pharm. Sci.*, 34, 187-194.
39. Viviane, G.C.A., Jacqueline, A.T., Lucienir, P.D., Dorila, P., Policarpo, A.S., Rosana, O.A., Alvaro, J.R., Antonio, F.C.A. (2011). Evaluation of the bactericidal and trypanocidal activities of triterpenes isolated from the leaves, stems and flowers of *Lychnophora pinaster*. *Brazilian journal of Pharmacognosy*, 21(4): 615-621.
40. Goyal, M.M., Gupta, A. (1987). Antibacterial activity of some active Principles of *polyathia longifolia* leaves, 19(3): 216-220.
41. Venkata, R., Samuel, L., Pardha, S.M., Narashimha, R. Naga, V.K.A., Sudhakar, M., and Radhakrishnan, T.M. (2012). Antibacterial, antioxidant activity and GC-MS Analysis of *Eupatorium odoratum*. *Asian J. Pharm. Clin. Res.*, 5, Suppl 2, 0974-2441.
42. Moorthy, V. Boominath, M. (2011). "The Antimicrobial Activities of Crude Extracts and fraction of *Psidium Guajava* and *Azadirachta indica* against *Staphylococcus aureus* in chronic disease affected patients". *International Journal of universal pharmacy and life sciences*, 1(2): September-October, 2011.