A Simple Mechanism for Gravitation

Chun-Xuan, Jiang (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄,北京 3924 信箱,100854)

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com, jcxxxx@163.com

Abstract: Gravity is a great mystery. No one has since given any machinery. In this paper we give a simple machinery. Gravity is the tachyon centripetal force. Anybody may understand gravitation.

[Chun-Xuan, Jiang (蒋春暄). A Simple Mechanism for Gravitation. Rep Opinion 2016;8(4):93-95]. ISSN 1553-9873 (print); ISSN 2375-7205 (online). http://www.sciencepub.net/report. 11. doi:10.7537/marsroj08041611.

Keywords: Gravity; mystery; machinery; tachyon; centripetal force.

Gravity is a great mystery. No one has since given any machinery. In this paper we give a simple machinery. Gravity is the tachyon centripetal force.

Anybody may understand gravitation.

Using the tardyon and tachyon coexistence principle [1]

$$u\overline{u} = c^2 \tag{1}$$

where c is light velocity in vacuum, $u \le c$ tardyon velocity and $\overline{u} \ge c$ tachyon velocity.

$$\overline{F} = -\frac{mc^2}{R}$$

We deduce the new gravitation formula:

Figure 1 shows that the rotation ω of body A emits tachyon mass \overline{m} , which forms the tachyon and gravitation field and gives the body B revolutions u and \overline{u} .




Fig. 1. On body $B = \frac{du}{dt}$ and $\frac{d\overline{u}}{dt}$ coexistence [2].

From Fig. 1 it follows

$$\frac{u\Delta t}{R} = \frac{\Delta u}{u} \tag{2}$$

From (2) it follows the tardyon centripetal acceleration on the body B [2-5],

$$\frac{du}{dt} = \lim_{\stackrel{\Delta u \to 0}{\Delta t \to 0}} \frac{\Delta u}{\Delta t} = \frac{u^2}{R}$$
(3)

From Fig. 1 it follows

$$\frac{u\Delta t}{R} = -\frac{\Delta \overline{u}}{\overline{u}} \tag{4}$$

From (4) and (1) it follows the tachyon centrifugal acceleration on the body $\,B\,$ [2-5],

$$\frac{d\overline{u}}{dt} = \lim_{\substack{\Delta \overline{u} \to 0 \\ \Delta t \to 0}} \frac{\Delta \overline{u}}{\Delta t} = -\frac{u\overline{u}}{R} = -\frac{c^2}{R}$$
(5)

On body $B = \frac{du}{dt}$ and $\frac{d\overline{u}}{dt}$ coexistence.

From (3) it follows the tardyon centrifugal force on body B [2-5],

$$F = \frac{M_B u^2}{R},\tag{6}$$

where M_B is body B mass.

From (5) it follows the tachyon centripetal force on body B, that is gravity [2-5],

$$\overline{F} = -\frac{mc^2}{R} \tag{7}$$

where m is the gravitation mass converted into by tachyon mass \overline{m} which is unobservable but m is observable. On body B F and \overline{F} coexistence.




Fig.2. On body B F and \overline{F} coexistence[2].

From Fig. 2, it follows

$$F + \overline{F} = 0. (8)$$

From (6), (7) and (8) it follows

$$\frac{m}{M_B} = \frac{u^2}{c^2} \tag{9}$$

Body B increases mass m and centrifugal force is greater than gravitation force, then body B expands outward. [5]

From (7) it follows Newtonian gravitation formula. The m is proportional to body A mass M_A , in (9) m is proportional to M_B , is inversely proportional to the distance R between body A and body B. It follows

$$m = k \frac{M_A M_B}{R} \tag{10}$$

where k is constant

Substituting (10) into (7) it follows the Newtonian gravitation formula [2-5]

$$\overline{F} = -G \frac{M_A M_B}{R^2} \tag{11}$$

where $G = kc^2 = 6.673 \times 10^{-8} \text{ cm}^3 / \text{g} \cdot \text{sec}^2$ is gravitation constant.

References

- 1. Chun-Xuan Jiang, A theory of morphisms between the tardyon and tachyon, Wuli(physics), (Chinese), 4. (175)119-125.
- 2. Chun-Xuan Jiang, On nature for gravitation, J. Beijing observatory (Chinese), 7(1976)32-38.
- 3. Chun-Xuan Jiang, An approach on the nature of attractive force, Potential science (Chinese), 4(1982)19-20.
- 4. Chun-Xuan Jiang, A unified theory of the gravitational and strong interactions, Hadronic J., 24(2001)629-638.

4/23/2016