

In 1991 Fermat's Last Theorem Has Been Proved(II)

Jiang, Chun-Xuan

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: In 1637 Fermat wrote: “It is impossible to separate a cube into two cubes, or a biquadrate into two biquadrates, or in general any power higher than the second into powers of like degree: I have discovered a truly marvelous proof, which this margin is too small to contain.” This means: $x^n + y^n = z^n$ ($n > 2$) has no integer solutions, all different from 0(i.e., it has only the trivial solution, where one of the integers is equal to 0). It has been called Fermat's last theorem (FLT). It suffices to prove FLT for exponent 4 and every prime exponent P . Fermat proved FLT for exponent 4. Euler proved FLT for exponent 3. In this paper using automorphic functions we prove FLT for exponents $4P$ and P , where P is an odd prime. We rediscover the Fermat proof. The proof of FLT must be direct. But indirect proof of FLT is disbelieving. In 1974 Jiang found out Euler formula of the cyclotomic

$$\exp\left(\sum_{i=1}^{4m-1} t_i J^i\right) = \sum_{i=1}^{4m} S_i J^{i-1}, \quad (1)$$

real numbers in the cyclotomic fields

where J denotes a $4m$ th root of unity, $J^{4m} = 1$, $m=1,2,3,\dots$, t_i are the real numbers.

S_i is called the automorphic functions(complex hyperbolic functions) of order $4m$ with $4m-1$ variables [2,5,7].

$$S_i = \frac{1}{4m} \left[e^{A_1} + 2e^H \cos\left(\beta + \frac{(i-1)\pi}{2}\right) + 2 \sum_{j=1}^{m-1} e^{B_j} \cos\left(\theta_j + \frac{(i-1)j\pi}{2m}\right) \right] \\ + \frac{(-1)^{(i-1)}}{4m} \left[e^{A_2} + 2 \sum_{j=1}^{m-1} e^{D_j} \cos\left(\phi_j - \frac{(i-1)j\pi}{2m}\right) \right] \quad (2)$$

where $i = 1, \dots, 4m$,

2000 Mathematics subject classification: 11D41

[Jiang, Chun-Xuan. **In 1991 Fermat's Last Theorem Has Been Proved(II).** *Researcher* 2016;8(8):21-24]. ISSN 1553-9865 (print); ISSN 2163-8950 (online). <http://www.sciencepub.net/researcher>. 5. doi:[10.7537/marsrj080816.05](https://doi.org/10.7537/marsrj080816.05).

Key words: complex hyperbolic, Fermat proof.

$$A_1 = \sum_{\alpha=1}^{4m-1} t_\alpha, \quad A_2 = \sum_{\alpha=1}^{4m-1} t_\alpha (-1)^\alpha, \quad H = \sum_{\alpha=1}^{2m-1} t_{2\alpha} (-1)^\alpha, \quad \beta = \sum_{\alpha=1}^{2m} t_{2\alpha-1} (-1)^\alpha, \\ B_j = \sum_{\alpha=1}^{4m-1} t_\alpha \cos \frac{\alpha j \pi}{2m}, \quad \theta_j = - \sum_{\alpha=1}^{4m-1} t_\alpha \sin \frac{\alpha j \pi}{2m}, \\ D_j = \sum_{\alpha=1}^{4m-1} t_\alpha (-1)^\alpha \cos \frac{\alpha j \pi}{2m}, \quad \phi_j = \sum_{\alpha=1}^{4m-1} t_\alpha (-1)^\alpha \sin \frac{\alpha j \pi}{2m}, \\ A_1 + A_2 + 2H + 2 \sum_{j=1}^{m-1} (B_j + D_j) = 0 \quad (3)$$

From (2) we have its inverse transformation[5,7]

$$\begin{aligned}
e^{A_1} &= \sum_{i=1}^{4m} S_i, \quad e^{A_2} = \sum_{i=1}^{4m} S_i (-1)^{1+i} \\
e^H \cos \beta &= \sum_{i=1}^{2m} S_{2i-1} (-1)^{1+i}, \quad e^H \sin \beta = \sum_{i=1}^{2m} S_{2i} (-1)^i \\
e^{B_j} \cos \theta_j &= S_1 + \sum_{i=1}^{4m-1} S_{1+i} \cos \frac{ij\pi}{2m}, \quad e^{B_j} \sin \theta_j = - \sum_{i=1}^{4m-1} S_{1+i} \sin \frac{ij\pi}{2m} \\
e^{D_j} \cos \phi_j &= S_1 + \sum_{i=1}^{4m-1} S_{1+i} (-1)^i \cos \frac{ij\pi}{2m}, \quad e^{D_j} \sin \phi_j = \sum_{i=1}^{4m-1} S_{1+i} (-1)^i \sin \frac{ij\pi}{2m}
\end{aligned} \tag{4}$$

(3) and (4) have the same form.

From (3) we have

$$\exp \left[A_1 + A_2 + 2H + 2 \sum_{j=1}^{m-1} (B_j + D_j) \right] = 1 \tag{5}$$

From (4) we have

$$\begin{aligned}
&\exp \left[A_1 + A_2 + 2H + 2 \sum_{j=1}^{m-1} (B_j + D_j) \right] = \begin{vmatrix} S_1 & S_{4m} & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \cdots \\ S_{4m} & S_{4m-1} & \cdots & S_1 \end{vmatrix} \\
&= \begin{vmatrix} S_1 & (S_1)_1 & \cdots & (S_1)_{4m-1} \\ S_2 & (S_2)_1 & \cdots & (S_2)_{4m-1} \\ \cdots & \cdots & \cdots & \cdots \\ S_{4m} & (S_{4m})_1 & \cdots & (S_{4m})_{4m-1} \end{vmatrix} \tag{6}
\end{aligned}$$

where

$$(S_i)_j = \frac{\partial S_i}{\partial t_j} [7]$$

From (5) and (6) we have circulant determinant

$$\exp \left[A_1 + A_2 + 2H + 2 \sum_{j=1}^{m-1} (B_j + D_j) \right] = \begin{vmatrix} S_1 & S_{4m} & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \cdots \\ S_{4m} & S_{4m-1} & \cdots & S_1 \end{vmatrix} = 1 \tag{7}$$

Assume $S_1 \neq 0, S_2 \neq 0, S_i = 0$, where $i = 3, \dots, 4m$. $S_i = 0$ are $(4m-2)$ indeterminate equations with $(4m-1)$ variables. From (4) we have

$$\begin{aligned}
e^{A_1} &= S_1 + S_2, \quad e^{A_2} = S_1 - S_2, \quad e^{2H} = S_1^2 + S_2^2 \\
e^{2B_j} &= S_1^2 + S_2^2 + 2S_1 S_2 \cos \frac{j\pi}{2m}, \quad e^{2D_j} = S_1^2 + S_2^2 - 2S_1 S_2 \cos \frac{j\pi}{2m}
\end{aligned} \tag{8}$$

Example [2]. Let $4m = 12$. From (3) we have

$$A_1 = (t_1 + t_{11}) + (t_2 + t_{10}) + (t_3 + t_9) + (t_4 + t_8) + (t_5 + t_7) + t_6,$$

$$\begin{aligned}
A_2 &= -(t_1 + t_{11}) + (t_2 + t_{10}) - (t_3 + t_9) + (t_4 + t_8) - (t_5 + t_7) + t_6, \\
H &= -(t_2 + t_{10}) + (t_4 + t_8) - t_6, \\
B_1 &= (t_1 + t_{11}) \cos \frac{\pi}{6} + (t_2 + t_{10}) \cos \frac{2\pi}{6} + (t_3 + t_9) \cos \frac{3\pi}{6} + (t_4 + t_8) \cos \frac{4\pi}{6} + (t_5 + t_7) \cos \frac{5\pi}{6} - t_6, \\
B_2 &= (t_1 + t_{11}) \cos \frac{2\pi}{6} + (t_2 + t_{10}) \cos \frac{4\pi}{6} + (t_3 + t_9) \cos \frac{6\pi}{6} + (t_4 + t_8) \cos \frac{8\pi}{6} + (t_5 + t_7) \cos \frac{10\pi}{6} + t_6, \\
D_1 &= -(t_1 + t_{11}) \cos \frac{\pi}{6} + (t_2 + t_{10}) \cos \frac{2\pi}{6} - (t_3 + t_9) \cos \frac{3\pi}{6} + (t_4 + t_8) \cos \frac{4\pi}{6} - (t_5 + t_7) \cos \frac{5\pi}{6} - t_6, \\
D_2 &= -(t_1 + t_{11}) \cos \frac{2\pi}{6} + (t_2 + t_{10}) \cos \frac{4\pi}{6} - (t_3 + t_9) \cos \frac{6\pi}{6} + (t_4 + t_8) \cos \frac{8\pi}{6} - (t_5 + t_7) \cos \frac{10\pi}{6} + t_6, \\
A_1 + A_2 + 2(H + B_1 + B_2 + D_1 + D_2) &= 0, \quad A_2 + 2B_2 = 3(-t_3 + t_6 - t_9). \tag{9}
\end{aligned}$$

From (8) and (9) we have

$$\exp[A_1 + A_2 + 2(H + B_1 + B_2 + D_1 + D_2)] = S_1^{12} - S_2^{12} = (S_1^3)^4 - (S_2^3)^4 = 1. \tag{10}$$

From (9) we have

$$\exp(A_2 + 2B_2) = [\exp(-t_3 + t_6 - t_9)]^3. \tag{11}$$

From (8) we have

$$\exp(A_2 + 2B_2) = (S_1 - S_2)(S_1^2 + S_2^2 + S_1 S_2) = S_1^3 - S_2^3. \tag{12}$$

From (11) and (12) we have Fermat's equation

$$\exp(A_2 + 2B_2) = S_1^3 - S_2^3 = [\exp(-t_3 + t_6 - t_9)]^3. \tag{13}$$

Fermat proved that (10) has no rational solutions for exponent 4 [8].

Therefore we prove we prove that (13) has no rational solutions for exponent 3. [2]

Theorem. Let $4m = 4P$, where P is an odd prime, $(P-1)/2$ is an even number.

From (3) and (8) we have

$$\exp[A_1 + A_2 + 2H + 2 \sum_{j=1}^{P-1} (B_j + D_j)] = S_1^{4P} - S_2^{4P} = (S_1^P)^4 - (S_2^P)^4 = 1 \tag{14}$$

From (3) we have

$$\exp[A_2 + 2 \sum_{j=1}^{\frac{P-1}{4}} (B_{4j-2} + D_{4j})] = [\exp(-t_p + t_{2p} - t_{3p})]^p. \tag{15}$$

From (8) we have

$$\exp[A_2 + 2 \sum_{j=1}^{\frac{P-1}{4}} (B_{4j-2} + D_{4j})] = S_1^P - S_2^P. \tag{16}$$

From (15) and (16) we have Fermat's equation

$$\exp[A_2 + 2 \sum_{j=1}^{\frac{P-1}{4}} (B_{4j-2} + D_{4j})] = S_1^P - S_2^P = [\exp(-t_p + t_{2p} - t_{3p})]^p \tag{17}$$

Fermat proved that (14) has no rational solutions for exponent 4 [8]. Therefor we prove that (17) has no rational solutions for prime exponent P .

Remark. Mathematicians said Fermat could not

possibly had a proof, because they do not understand FLT. In complex hyperbolic functions let exponent n be $n = \Pi P$, $n = 2\Pi P$ and $n = 4\Pi P$. Every factor of exponent n has Fermat's equation [1-7]. Using modular elliptic curves Wiles and Taylor prove

FLT [9,10]. They has not proved FLT[11,12] The classical theory of automorphic functions, created by Klein and Poincare, was concerned with the study of analytic functions in the unit circle that are invariant under a discrete group of transformation. Automorphic functions are the generalization of trigonometric, hyperbolic elliptic, and certain other functions of elementary analysis. The complex trigonometric functions and complex hyperbolic functions have a wide application in mathematics and physics.

References

1. Jiang, C-X, Fermat last theorem had been proved, Potential Science (in Chinese), 2.17-20 (1992), Preprints (in English) December (1991). <http://www.wbabin.net/math/xuan47.pdf>.
2. Jiang, C-X, Fermat last theorem had been proved by Fermat more than 300 years ago, Potential Science (in Chinese), 6.18-20(1992).
3. Jiang, C-X, On the factorization theorem of circulant determinant, Algebras, Groups and Geometries, 11. 371-377(1994), MR. 96a: 11023, <http://www.wbabin.net/math/xuan45.pdf>
4. Jiang, C-X, Fermat last theorem was proved in 1991, Preprints (1993). In: Fundamental open problems in science at the end of the millennium, T. Gill, K. Liu and E. Trell (eds). Hadronic Press, 1999, P555-558. <http://www.wbabin.net/math/xuan46.pdf>.
5. Jiang, C-X, On the Fermat-Santilli theorem, Algebras, Groups and Geometries, 15. 319-349(1998).
6. Jiang, C-X, Complex hyperbolic functions and Fermat's last theorem, Hadronic Journal Supplement, 15. 341-348(2000).
7. Jiang, C-X, Foundations of Santilli Isonumber Theory with applications to new cryptograms, Fermat's theorem and Goldbach's Conjecture. Inter. Acad. Press. 2002. MR2004c:11001, <http://www.wbabin.net/math/xuan13.pdf>. <http://www.i-b-r.org/docs/jiang.pdf>.
8. Ribenboim, P, Fermat last theorem for amateur, Springer-Verlag, (1999).
9. Wiles A, Modular elliptic curves and Fenmat's last theorem, Ann of Math, (2) 141 (1995), 443-551.
10. Taylor, R. and Wiles, A., Ring-theoretic properties of certain Hecke algebras, Ann. of Math., (2) 141(1995), 553-572.
11. Zhivolov, Y, Fermat last theorem and mistakes of Andrew Wiles, (2006). www.baoway.com/bbs/viewthread.php?tid=18162&fpage=3.
12. Zhivolov, Y, Fermat last theorem and Kenneth Ribet mistakes, (2006). www.baoway.com/bbs/viewthread.php?tid=18164&fpage=3.
13. Jiang, Chun-Xuan (蒋春暄). Automorphic Functions And Fermat's Last Theorem (1). *Academ Arena* 2016;8(5):78-84]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 8. doi:[10.7537/marsaa080516.08](https://doi.org/10.7537/marsaa080516.08).
14. Jiang, Chun-Xuan (蒋春暄). Automorphic Functions And Fermat's Last Theorem (3) (Fermat's Proof of FLT). *Academ Arena* 2016;8(5):85-92]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 9. doi:[10.7537/marsaa080516.09](https://doi.org/10.7537/marsaa080516.09).
15. Jiang, Chun-Xuan (蒋春暄). Riemann Paper (1859) Is False. *Academ Arena* 2016;8(6):35-39. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 7. doi:[10.7537/marsaa080616.07](https://doi.org/10.7537/marsaa080616.07).
16. Jiang, Chun-Xuan (蒋春暄). The New Prime theorems (341) - (390) . *Academ Arena* 2016;8(6):61-107]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 12. doi:[10.7537/marsaa080616.12](https://doi.org/10.7537/marsaa080616.12).
17. Baidu. <http://www.baidu.com>. 2016.
18. Google. <http://www.google.com>. 2016.
19. Ma H, Yang Y. Turritopsis nutricula. Nature and Science 2010;8(2):15-20. doi:[10.7537/marsnsj080210.03](https://doi.org/10.7537/marsnsj080210.03). http://www.sciencepub.net/nature/ns0802/03_1279_hongbao_turritopsis_ns0802_15_20.pdf.
20. Ma H. The Nature of Time and Space. Nature and science 2003;1(1):1-11. doi:[10.7537/marsnsj010103.01](https://doi.org/10.7537/marsnsj010103.01). <http://www.sciencepub.net/nature/0101/01-ma.pdf>.
21. Wikipedia. The free encyclopedia. <http://en.wikipedia.org>. 2016.
22. 马宏宝. 宙永恒吗? New York Science Journal. 2008;1(3):66-69. ISSN: 1554-0200. doi:[10.7537/marsnys010308.07](https://doi.org/10.7537/marsnys010308.07). http://www.sciencepub.net/newyork/0103/07_0026_mahongbao_universe.pdf.